

THIN Bibliography

| No. | Researcher | Title | Journal | Reference | Year |
|-----|---|---|---|--|------|
| 545 | Lycett D, Nichols L, Ryan R, Farley A, Roalfe A, Mohammed MA, Szatkowski L, Coleman T, Morris R, Farmer A & Aveyard P | The association between smoking cessation and glycaemic control in patients with type 2 diabetes: a THIN database cohort study | Lancet Diabetes Endocrinol | 3(6):423-30 | 2015 |
| 544 | Boursi B, Mamtani R, Haynes K & Yang YX | The effect of past antibiotic exposure on diabetes risk | Eur J Endocrinol | 172(6):639-48 | 2015 |
| 543 | O'Leary C, Nasser A, Myland M, Waples S & Ansell D | Audit of irreversible monoamine oxidase inhibitor (MAOI) prescribing for depression in current clinical practice using The Health Improvement Network (THIN) UK primary care database | ISPOR 20th Annual International meeting | Philadelphia, USA | 2015 |
| 542 | Moran GM, Calvert M, Feltham MG, Ryan R & Marshall T | A retrospective cohort study to investigate fatigue, psychological or cognitive impairment after TIA: protocol paper | BMJ Open | 3;5(4):e008149 | 2015 |
| 541 | Toulis KA, Nirantharakumar K, Ryan R, Marshall T & Hemming K | Bisphosphonates and glucose homeostasis: a population-based, retrospective cohort study | J Clin Endocrinol Metab | 100(5):1933-40 | 2015 |
| 540 | Star K, Watason S, Sandberg L, Johansson J, Edwards IR | Longitudinal medical records as a complement to routine drug safety signal analysis | Pharmacoepidemiol Drug Saf | 24(5):486-94 | 2015 |
| 539 | Zingone F, West J, Crooks CJ, Fleming KM, Card TR, Ciacci C & Tata LJ | Socioeconomic variation in the incidence of childhood coeliac disease in the UK | Arch Dis Child | 100(5):466-73 | 2015 |
| 538 | O'Leary C, Anderson P, Junor R, Marsden D, Nasser A, Pockett R & Ansell D | HECTOR Study – Health Economic study of Codeine and Tramadol nOn-Responding patients with chronic non-malignant pain using The Health Improvement Network (THIN) UK primary care database: an interim analysis | The British Pain Society 2015 Annual Scientific Meeting (ASM) | Glasgow at the SECC: Scottish Exhibition and Conference Centre | 2015 |
| 537 | Saine ME, Carbonari DM, Newcomb CW, et al. | Determinants of saxagliptin use among patients with type 2 diabetes mellitus treated with oral anti-diabetic drugs | BMC Pharmacol toxicol | 16(1):8 | 2015 |
| 536 | Raluy-Callado M, Gabriel S, Dinet J, Wang M & Wasiaik R | A retrospective study to assess resource utilization in patients with cervical dystonia in the United Kingdom | Neuropsychiatr Dis Treat | 9;11:647-55 | 2015 |
| 535 | Meeraus WH, Petersen I, Gilbert R | Association between Antibiotic Prescribing in Pregnancy and Cerebral Palsy or Epilepsy in Children Born at Term: A Cohort Study Using The Health Improvement Network | PLoS One | 10(3):e0122034 | 2015 |
| 534 | Osborn DP, Hardoon S, Omar RZ, Holt RI, et al. | Cardiovascular risk prediction models for people with severe mental illness: results from the prediction and management of cardiovascular risk in people with severe mental illnesses (PRIMROSE) research program | JAMA Psychiatry | 72(2):143-51 | 2015 |
| 533 | Michel A, Martin-Perez M, Ruigomez A, Garcia Rodriguez LA | Risk factors for hyperkalaemia in a cohort of patients with newly diagnosed heart failure: a nested case-control study in UK general practice | Eur J Heart Fail | 17(2):205-13 | 2015 |

| | | | | | |
|-----|---|--|--|------------------|------|
| 532 | O'Leary C, Nasser A, Myland M, Waples S & Ansell D | Audit of irreversible monoamine oxidase inhibitor (MAOI) prescribing for depression in current clinical practice using The Health Improvement Network (THIN) UK primary care database | PRIMM conference | London, UK | 2015 |
| 531 | Nasser A, O'Leary C, Myland M, Sam Waples S & Ansell D | Audit of irreversible monoamine oxidase inhibitors (MAOI) prescription for depression in current clinical practice within The Health Improvement Network (THIN) UK primary care database | PRIMM conference, 26th Annual Scientific Meeting, Imperial Hotel, Russell Square, London | [Ahead of print] | 2015 |
| 530 | Masclee GM, Coloma PM, Spaander MC, Kuipers EJ & Sturkenboom MC | NSAIDs, statins, low-dose aspirin and PPIs, and the risk of oesophageal adenocarcinoma among patients with Barrett's oesophagus: a population-based case-control study | BMJ Open | 29;5(1):e006640 | 2015 |
| 529 | Schrag A, Horsfall L, Walters K, Noyce A & Petersen I | Prediagnostic presentations of Parkinson's disease in primary care: a case-control study | The Lancet | 14(1):57-64 | 2015 |
| 528 | Cederholm S, Hill G, Asimwe A, Bate A, Bhayat F, Persson Brobert G, Bergvall T, Ansell D, Star K & Norén GN | Structured assessment for prospective identification of safety signals in electronic medical records: evaluation in the health improvement network | Drug Saf | 38(1):87-100 | 2015 |
| 527 | Shallcross LJ, Hayward AC, Johnson AM, Petersen I | Evidence for increasing severity of community-onset boils and abscesses in UK General Practice | Epidemiol Infect | 22:1-4 | 2014 |
| 526 | Keane MG, Horsfall L, Rait G & Pereira SP | A case-control study comparing the incidence of early symptoms in pancreatic and biliary tract cancer | BMJ Open | 19;4(11):e005720 | 2014 |
| 525 | Keane MG, Horsfall L, Rait G & Pereira SP | A case-control study comparing the incidence of early symptoms in pancreatic and biliary tract cancer | BMJ Open | 19;4(11):e005720 | 2014 |
| 524 | Dhalwani NN, Tata LJ, Coleman T, Fiaschi L & Szatkowski L | A comparison of UK primary care data with other national data sources for monitoring the prevalence of smoking during pregnancy | J Public Health (Oxf) | [ahead of print] | 2014 |
| 523 | Buszewicz M, Welch C, Horsfall L, Nazareth I, Osborn D, et al. | Assessment of an incentivised scheme to provide annual health checks in primary care for adults with intellectual disability: a longitudinal cohort study | The Lancet | [Ahead of print] | 2014 |
| 522 | Boursi B, Haynes K, Mamtani R & Yang YX | Digoxin use and the risk for colorectal cancer | Pharmacoepidemiol Drug Saf | 23(11):1147-53 | 2014 |
| 521 | Carey IM, Shah SM, Dewilde S & et al | Increased Risk of Acute Cardiovascular Events After Partner Bereavement: A Matched Cohort Study | JAMA Intern Med | 174(4):598-605 | 2014 |
| 520 | Szatkowski L, Murray R, Hubbard R, Agrawal S, Huang Y & Britton J | Prevalence of smoking among patients treated in NHS hospitals in England in 2010/2011: a national audit | Thorax | [ahead of print] | 2014 |
| 519 | McEwan P, Bennett H, Ward T & Bergenheim K | Refitting of the UKPDS 68 Risk Equations to Contemporary Routine Clinical Practice Data in the UK | Pharmacoeconomics | [ahead of print] | 2014 |
| 518 | Moran GM, Calvert M, Feltham MG & Marshall T | Retrospective case review of missed opportunities for primary prevention of stroke and TIA in primary care: protocol paper | BMJ Open | 11;4(11):e006622 | 2014 |
| 517 | Baker R, Orton E, Tata LJ & Kendrick D | Risk factors for long-bone fractures in children up to 5 years of age: a nested case-control study | Arch Dis Child | [ahead of print] | 2014 |
| 516 | Ogdie A, Yu Y, Haynes K, Love TJ, Maliha S, Jiang Y, Troxel AB, Hennessy S, Kimmel SE, Margolis DJ, Choi H, Mehta NN & Gelfand JM | Risk of major cardiovascular events in patients with psoriatic arthritis, psoriasis and rheumatoid arthritis: a population-based cohort study | Ann Rheum Dis | [ahead of print] | 2014 |
| 515 | Mamtani R, Haynes K, Boursi B, Scott F, Goldberg DS, Keefe SM, Vaughn DJ, Malkowicz SB & Lewis JD | Validation of a coding algorithm to identify bladder cancer and distinguish stage in an electronic medical records database | Cancer Epidemiol Biomarkers Prev | [Ahead of print] | 2014 |

| | | | | | |
|-----|--|--|---|-----------------------|------|
| 514 | Petersen I, McCrea RL, Osborn DJ, Evans S, Pinfold V, Cowen PJ, Gilbert R & Nazareth I | Discontinuation of antipsychotic medication in pregnancy: a cohort study | Schizophr Res | 159(1):218-25 | 2014 |
| 513 | Hakimi Z, Johnson M, Nazir J, Blak B & Odeyemi IA | Drug treatment patterns for the management of men with lower urinary tract symptoms associated with benign prostatic hyperplasia who have both storage and voiding symptoms: a study using the health improvement network UK primary care data | Curr Med Res Opin | [ahead of print] | 2014 |
| 512 | Takeshita J, Wang S, Shin DB, Mehta NN, Kimmel SE, Margolis DJ, Troxel AB & Gelfand JM | Effect of Psoriasis Severity on Hypertension Control: A Population-Based Study in the United Kingdom | JAMA Dermatol | [ahead of print] | 2014 |
| 511 | Rho YH, Lu N, Peloquin CE, Man A, Zhu Y, Zhang Y & Choi HK | Independent impact of gout on the risk of diabetes mellitus among women and men: a population-based, BMI-matched cohort study | Ann Rheum Dis | [ahead of print] | 2014 |
| 510 | Cea Soriano L, Bateman BT, García Rodríguez LA & Hernández-Díaz S | Prescription of antihypertensive medications during pregnancy in the UK | Pharmacoepidemiol Drug Saf | 23(10):1051-8 | 2014 |
| 509 | Huang Y, Lewis S & Britton J | Use of varenicline for smoking cessation treatment in UK primary care: an association rule mining analysis | BMC Public Health | 2;14:1024 | 2014 |
| 508 | Bennett H, McEwan P, Bergenheim K & Gordon J | Assessment of Unmet Clinical Need in Type 2 Diabetic Patients on Conventional Therapy in the UK | Diabetes Ther | [ahead of print] | 2014 |
| 507 | Walker AM, Vannappagari V, Buus R, Covington D, Bratton E & Thompson M | Descriptive Study of Exposure to Inhalation Zanamivir and Pregnancy-Related Outcomes | Journal of Pharmaceutics & Drug Development | 1(5), ISSN: 2348-9782 | 2014 |
| 506 | Dood RL, Gracia CR, Sammel MD, Haynes K, Senapati S, Strom BL | Endometrial cancer after endometrial ablation vs medical management of abnormal uterine bleeding | J Minim Invasive Gynecol | 21(5):744-52 | 2014 |
| 505 | Welch CA, Petersen I, Bartlett JW, White IR, Marston L, Morris RW, Nazareth I, Walters K & Carpenter J | Evaluation of two-fold fully conditional specification multiple imputation for longitudinal electronic health record data. | Statistics in Medicine | 20;33(21):3725-37 | 2014 |
| 504 | O'Keeffe AG, Geneletti S, Baio G, Sharples LD, Nazareth I & Petersen I | Regression discontinuity designs: an approach to the evaluation of treatment efficacy in primary care using observational data | BMJ | 8;349:g5293 | 2014 |
| 503 | Keane MG, Horsfall LJ Rait G2 & Pereira SP | Sociodemographic trends in the incidence of pancreatic and biliary tract cancer in UK primary care | PLoS One | 30;9(9):e108498 | 2014 |
| 502 | Cea Soriano L, Wallander MA, Andersson S, Filonenko A & García Rodríguez LA | The continuation rates of long-acting reversible contraceptives in UK general practice using data from The Health Improvement Network | Pharmacoepidemiol Drug Saf | [ahead of print] | 2014 |
| 501 | Ogdie A, Alehashemi S, Love TJ, Jiang Y, Haynes K, Hennessy S, Choi H & Gelfand JM | Validity of psoriatic arthritis and capture of disease modifying antirheumatic drugs in the health improvement network | Pharmacoepidemiol Drug Saf | 23(9):918-22 | 2014 |
| 500 | Lind M, Pivodic A, Cea-Soriano L, Nerman O, Pehrsson NG & Garcia-Rodriguez LA | Changes in HbA1c and frequency of measuring HbA1c and adjusting glucose-lowering medications in the 10 years following diagnosis of type 2 diabetes: a population-based study in the UK | Diabetologia | 57(8):1586-94 | 2014 |
| 499 | Sáez ME, González-Pérez A, Johansson S, Nagy P & García Rodríguez LA | Patterns in the Use of Low-Dose Acetylsalicylic Acid and Other Therapies Following Upper Gastrointestinal Bleeding | American Journal of Cardiovascular Drugs | [ahead of print] | 2014 |

| | | | | | |
|-----|--|--|--|------------------------------------|------|
| 498 | Martín-Merino E, Johansson S, Nagy P & Rodríguez LA | Represcription of Low-Dose Acetylsalicylic Acid After Discontinuation in Patients Receiving Treatment for Secondary Cardiovascular Disease Prevention in the UK | American Journal of Cardiovascular Drugs | 14(4):319-26 | 2014 |
| 497 | Jain P, Calvert M, Cockwell P & McManus RJ | The need for improved identification and accurate classification of stages 3-5 Chronic Kidney Disease in primary care: retrospective cohort study | PLoS One | 12;9(8):e100831 | 2014 |
| 496 | Jones MP, Walker MM, Ford AC & Talley NJ | The overlap of atopy and functional gastrointestinal disorders among 23,471 patients in primary care | Aliment Pharmacol Ther | 40(4):382-91 | 2014 |
| 495 | Cea Soriano L, Wallander MA, Andersson S, Filonenko A & García Rodríguez LA | Use of long-acting reversible contraceptives in the UK from 2004 to 2010: Analysis using The Health Improvement Network Database | Eur J Contracept Reprod Health Care | 19(6):439-47 | 2014 |
| 494 | Cea Soriano L, Wallander MA, Andersson S, Filonenko A & García Rodríguez LA | Use of long-acting reversible contraceptives in the UK from 2004 to 2010: Analysis using The Health Improvement Network Database | The European Journal of Contraception and Reproductive Health Care | [ahead of print] | 2014 |
| 493 | Breuer J, Pacou M, Gautier A & Brown MM | Herpes zoster as a risk factor for stroke and TIA: a retrospective cohort study in the UK | Neurology | 8;83(2):e27-33 | 2014 |
| 492 | Misra D, Zhang Y, Peloquin C, Choi HK, Kiel DP & Neogi T | Incident long-term warfarin use and risk of osteoporotic fractures: propensity-score matched cohort of elders with new onset atrial fibrillation | Osteoporosis International | 25(6):1677-84 | 2014 |
| 491 | González-Pérez A, Sáez ME, Johansson S, Nagy P & García Rodríguez LA | Risk factors associated with uncomplicated peptic ulcer and changes in medication use after diagnosis | PLoS One | 8;9(7):e101768 | 2014 |
| 490 | Ogdie A, Alehashemi S, Love TJ, Jiang Y, Haynes K, Hennessy S, Choi H & Gelfand JM | Validity of psoriatic arthritis and capture of disease modifying antirheumatic drugs in the health improvement network | Pharmacoepidemiology and Drug Safety | doi: 10.1002/pds.3677 | 2014 |
| 489 | O'Leary C, Bourke A & Ansell D | Comparing The Influence Of Month Of Birth And Gender In Two Academic Years On Attention Deficit Hyperactivity Diagnoses (ADHD) Among Children In The Health Improvement Network (THIN) UK Data | 19th Annual International Meeting (ISPOR) | Montreal, Canada | 2014 |
| 488 | Hall GC & Hill F | Descriptive investigation of the recording of influenza vaccination details on The Health Information Network database | Pharmacoepidemiology and Drug Safety | 23(6):595-600 | 2014 |
| 487 | Groves-Kirkby CJ, Denman AR, Campbell J, Rogers S, Phillips PS & Crockett RGM | Does Environmental Radon Gas Influence the Incidence of Multiple Sclerosis? A Preliminary Retrospective Study in England and Wales | The European IRPA Congress | Geneva | 2014 |
| 486 | Martín-Merino E, Fortuny J, Rivero-Ferrer Z & García-Rodríguez LA | Incidence of retinal complications in a cohort of newly diagnosed diabetic patients | PLoS One | 25;9(6):e100283 | 2014 |
| 485 | Misra D, Zhang Y, Peloquin C, Choi HK, Kiel DP & Neogi T | Incident long-term warfarin use and risk of osteoporotic fractures: propensity-score matched cohort of elders with new onset atrial fibrillation | Osteoporosis International | 25(6):1677-84 | 2014 |
| 484 | Ansell D, O'Leary C & Nasser A | Longitudinal Analysis Of Dengue Fever Infections Reported In The UK Between 2002 – 2013 Using The Health Improvement Network (Thin) Primary Care Database | 19th Annual International Meeting (ISPOR) | Montreal, Canada | 2014 |
| 483 | Horsfall LJ, Nazareth I & Petersen I | Serum uric acid and the risk of respiratory disease: a population-based cohort study | Thorax | doi: 10.1136/thoraxjnl-2014-205271 | 2014 |

| | | | | | |
|-----|---|--|---|----------------------------|------|
| 482 | James GD, Donaldson GC, Wedzicha JA & Nazareth I | Trends in management and outcomes of COPD patients in primary care, 2000-2009: a retrospective cohort study | NPJ Primary Care Respiratory Medicine | 3;24:14015 | 2014 |
| 481 | Karch A & Thomas SL | Autoimmune thyroiditis as a risk factor for stroke: A historical cohort study | Neurology | 82(18):1643-52 | 2014 |
| 480 | Guest JF, Panca M, Sladkevicius E, Taheri S & Stradling J | Clinical outcomes and cost-effectiveness of continuous positive airway pressure to manage obstructive sleep apnea in patients with type 2 diabetes in the U.K | Diabetes Care | 37(5):1263-71 | 2014 |
| 479 | Mamza J, Mehta R, Idris I | Obesity independently predicts responders to biphasic insulin 50/50 (Humalog Mix50 and Insuman Comb 50) following conversion from other insulin regimens: a retrospective cohort study | BMJ Open Diabetes Res Care | 7;2(1):e000021 | 2014 |
| 478 | Cea Soriano L, Bateman BT, García Rodríguez LA & Hernández-Díaz S | Prescription of antihypertensive medications during pregnancy in the UK | Pharmacoepidemiology and Drug Safety | [ahead of print] | 2014 |
| 477 | Masclee GMC, Coloma PM, de Wilde M, Kuipers EJ & Sturkenboom M | The incidence of Barrett's oesophagus and oesophageal adenocarcinoma in the United Kingdom and the Netherlands is levelling off | Alimentary Pharmacology and Therapeutics | doi:10.1111/apt.12759 | 2014 |
| 476 | Helbert M, Holbrook T, MacCulloch A & Mannan A | Understanding the cost of hereditary angioedema in England | The European Conference on Rare Diseases and Orphan Products (ECRD) | Berlin, Germany | 2014 |
| 475 | Shah P, Rhim AD, Haynes K, Hwang WT & Yang YX | Diagnosis of pernicious anemia and the risk of pancreatic cancer | Pancreas | 43(3):422-6 | 2014 |
| 474 | Cooper S, Menon S, Nightingale P & Trudgill N | Risk factors for the development of oesophageal adenocarcinoma in Barrett's oesophagus: a UK primary care retrospective nested case-control study | United European Gastroenterol J | 2(2):91-8 | 2014 |
| 473 | Eftekhari K, Ghodasra DH, Haynes K, Chen J, Kempen JH & Vanderbeek BL | Risk of retinal tear or detachment with oral fluoroquinolone use: a cohort study | Pharmacoepidemiology and Drug Safety | [ahead of print] | 2014 |
| 472 | Marston L & Carpenter JR, Walters KR, Morris RW, Nazareth I, White IR & Petersen I | Smoker, ex-smoker or non-smoker? The validity of routinely recorded smoking status in UK primary care: a cross-sectional study | BMJ Open | 23;4(4):e004958 | 2014 |
| 471 | Cooper S, Menon S, Nightingale P & Trudgill N | Risk factors for the development of oesophageal adenocarcinoma in Barrett's oesophagus: a UK primary care retrospective nested case-control study | United European Gastroenterol J | 2(2):91-8 | 2014 |
| 470 | Titchener AG, White JJ, Hinchliffe SR, Tamba AA, Hubbard RB & Clark D | Comorbidities in rotator cuff disease: a case-control study | Journal of Shoulder and Elbow Surgery | pii: S1058-2746(13)00644-7 | 2014 |
| 469 | Cea Soriano L, Ruigómez A, Johansson S & García Rodríguez LA | Study of the Association Between Hip Fracture and Acid-Suppressive Drug Use in a UK Primary Care Setting | Pharmacotherapy | [ahead of print] | 2014 |
| 468 | Dubreuil M, Rho YH, Man A, Zhu Y, Zhang Y, Love TJ, Ogdie A, Gelfand JM & Choi HK | Diabetes incidence in psoriatic arthritis, psoriasis and rheumatoid arthritis: a UK population-based cohort study | Rheumatology | 53(2):346-52 | 2014 |
| 467 | Mamtani R, Pfanzer N, Haynes K, Finkelman BS, Wang X, Keefe SM, Haas NB, Vaughn DJ & Lewis JD | Incidence of Bladder Cancer in Patients With Type 2 Diabetes Treated With Metformin or Sulfonylureas | Diabetes Care | [ahead of print] | 2014 |
| 466 | Hardy B, Szatkowski L, Tata LJ, Coleman T & Dhalwani NN | Smoking cessation advice recorded during pregnancy in United Kingdom primary care | BMC Family Practice | 1;15(1):21 | 2014 |

| | | | | | |
|-----|--|--|---|-----------------------|------|
| 465 | Donaldson JW, McKeever TM, Hall IP, Hubbard RB, Fogarty AW | The UK prevalence of hereditary haemorrhagic telangiectasia and its association with sex, socioeconomic status and region of residence: a population-based study | Thorax | 69(2):161-7 | 2014 |
| 464 | Dhiman P, Kai J, Horsfall L, Walters K & Qureshi N | Availability and quality of coronary heart disease family history in primary care medical records: implications for cardiovascular risk assessment | PLoS One | 9;9(1):e81998 | 2014 |
| 463 | Flory JH, Small DS, Cassano PA, Brillon DJ, Mushlin AI & Hennessy S | Comparative effectiveness of oral diabetes drug combinations in reducing glycosylated hemoglobin | Journal of comparative effectiveness research | 3(1):29-39 | 2014 |
| 462 | Fett N, Haynes K, Propert KJ & Margolis DJ | Five-year malignancy incidence in patients with chronic pruritus: A population-based cohort study aimed at limiting unnecessary screening practices | Journal of the American Academy of Dermatology | S0190-9622(13)01304-2 | 2014 |
| 461 | Breuer J, Pacou M, Gauthier A & Brown MM | Herpes zoster as a risk factor for stroke and TIA: A retrospective cohort study in the UK | Neurology | 21;82(3):206-12 | 2014 |
| 460 | Lin KJ, De Caterina R & Rodríguez LA | Low-dose aspirin and upper gastrointestinal bleeding in primary versus secondary cardiovascular prevention: a population-based, nested case-control study | Circulation: Cardiovascular Quality and Outcomes | 1;7(1):70-7 | 2014 |
| 459 | Mamtani R, Haynes K, Finkelman BS, Scott FI & Lewis JD | Distinguishing incident and prevalent diabetes in an electronic medical record database | Pharmacoepidemiology and Drug Safety | [ahead of print] | 2013 |
| 458 | Helbert M, Holbrook T, Drogon E & Mannan A | Exploring the cost and burden of illness of hereditary angioedema in the UK | UK PIN Liverpool | Liverpool, UK | 2013 |
| 457 | Shah SM, Carey IM, Harris T & et al | Impact of partner bereavement on quality of cardiovascular disease management | Circulation | 128(25):2745-2753 | 2013 |
| 456 | Shah SM, Carey IM, Harris T & et al | Initiation of psychotropic medication after partner bereavement: a matched cohort study | PLoS One | 8(11):e77734 | 2013 |
| 455 | Lind M, Garcia-Rodriguez LA, Booth GL, Ceasoriano L, Shah BR, Ekeroth G & Lipscombe LL | Mortality trends in patients with and without diabetes in Ontario, Canada and the UK from 1996 to 2009: a population-based study | Diabetologia | 56(12):2601-8 | 2013 |
| 454 | Hall GC, Amber V, O'Regan C & Jameson K | Observational study of ezetimibe discontinuation in primary care practices in the UK | Current Medical Research & Opinion | 29(12):1737-45 | 2013 |
| 453 | Hardoon S, Hayes JF, Blackburn R, Petersen I, Walters K, Nazareth I & Osborn D | Recording of severe mental illness in United kingdom primary care, 2000-2010 | Public Library of Science (PLoS one) | 12;8(12):e82365 | 2013 |
| 452 | Shah SM, Carey IM, Harris T & et al | The effect of unexpected bereavement on mortality in older couples | Am J Public Health | 103(6):1140-1145 | 2013 |
| 451 | Ray K, Foody J, Khan I, Gorcyca K, Ansell D & Lewis B | Attainment of Low-Density Lipoprotein Cholesterol Goals in Patients at Very High Cardiovascular Risk in the UK: Results from a General Practice Population Study | 16th Annual European Congress (ISPOR) | Dublin, Ireland | 2013 |
| 450 | Langley T, Lewis S, McNeill A, Gilmore A, Szatkowski L et al. | Characterizing tobacco control mass media campaigns in England | Addiction | 108(11):2001-8 | 2013 |
| 449 | Helbert M, Drogon E, Mannan A, Holbrook T, Murray H & Piercy J | Exploring the burden and cost of illness of hereditary angioedema in the UK | International Society for Pharmacoeconomics and Outcomes Research (ISPOR) 16th Annual European Congress | Dublin, Ireland | 2013 |

| | | | | | |
|-----|--|--|--|------------------|------|
| 448 | Valkhoff VE, Schade R, 't Jong GW, Romio S, et al | Population-based analysis of non-steroidal anti-inflammatory drug use among children in four European countries in the SOS project: what size of data platforms and which study designs do we need to assess safety issues | BMC Pediatr | 19;13:192 | 2013 |
| 447 | Cea Soriano L, Wallander MA, Andersson SW, Requena G & García-Rodríguez LA | Study of long-acting reversible contraceptive use in a UK primary care database: Validation of methodology | Eur J Contracept Reprod Health Care | [Ahead of print] | 2013 |
| 446 | Khadjesari Z, Marston L, Petersen I, Nazareth I & Walters K | Alcohol consumption screening of newly-registered patients in primary care: a cross-sectional analysis | British Journal of General Practice | 63(615):706-12 | 2013 |
| 445 | Hwang AL, Haynes K, Hwang WT, Yang YX | Metformin and survival in pancreatic cancer: a retrospective cohort study | Pancreas | 42(7):1054-9 | 2013 |
| 444 | Wan J, Wang S, Haynes K, Denburg MR, Shin DB & Gelfand JM | Risk of moderate to advanced kidney disease in patients with psoriasis: population based cohort study | British Medical Journal (BMJ) | 15;347:f5961 | 2013 |
| 443 | Cea-Soriano L, García Rodríguez L, Machlitt A & Wallander MA | Use of prescription contraceptive methods in the UK general population: a primary care study | An International Journal of Obstetrics & Gynaecology (BJOG) | [Ahead of print] | 2013 |
| 442 | Cea-Soriano L, García Rodríguez LA, Fernández Cantero O & Hernández-Díaz S | Challenges of using primary care electronic medical records in the UK to study medications in pregnancy | Pharmacoepidemiology and Drug Safety | 22(9):977-85 | 2013 |
| 441 | Dhalwani NN, Tata LJ, Coleman T, Fleming KM & Szatkowski L | Completeness of maternal smoking status recording during pregnancy in United Kingdom primary care data | Public Library of Science (PLoS one) | 19;8(9):e72218 | 2013 |
| 440 | Wijlaars LP, Nazareth I, Whitaker HJ, Evans SJ & Petersen I | Suicide-related events in young people following prescription of SSRIs and other antidepressants: a self-controlled case series analysis | BMJ Open | 19;3(9):e003247 | 2013 |
| 439 | Johnson M, Vial D, Thompson M, Cockbain C, Hogan S & Bourke A | A new method to get fast patient reported outcomes data from primary care in the UK for patients with atrial fibrillation (AF) in The Health Improvement Network (THIN) | 29th International Conference on Pharmacoepidemiology and Therapeutic Risk Management (ICPE) | Montreal, Canada | 2013 |
| 438 | García-Rodríguez LA, Gaist D, Morton J, Cookson C & González-Pérez A | Antithrombotic drugs and risk of hemorrhagic stroke in the general population | Neurology | 6;81(6):566-74 | 2013 |
| 437 | Johnson M, Mook P, Thompson M & Bourke A | Audit of the effect of relative age on morbidity among children in The Health Improvement Network (THIN) UK data | 29th International Conference on Pharmacoepidemiology and Therapeutic Risk Management (ICPE) | Montreal, Canada | 2013 |
| 436 | Johnson M, Thompson M, Mook P, Hogan S & Bourke A | Cause of death recording in The Health Improvement Network (THIN) UK data among the AFLOAT study population (N=56,508) | 29th International Conference on Pharmacoepidemiology and Therapeutic Risk Management (ICPE) | Montreal, Canada | 2013 |
| 435 | Grant RL, Drennan VM, Rait G, Petersen I & Iliffe S | First diagnosis and management of incontinence in older people with and without dementia in primary care: a cohort study using The Health Improvement Network primary care database | Public Library of Science (PLoS one) | 10(8):e1001505 | 2013 |
| 434 | McKeever T, Harrison TW, Hubbard R & Shaw D | Inhaled Corticosteroids and the Risk of Pneumonia in people with Asthma: A case control study | Chest | [ahead of print] | 2013 |
| 433 | González-Pérez A, Gaist D, Wallander MA, McFeat G & García-Rodríguez LA | Mortality after hemorrhagic stroke: Data from general practice (The Health Improvement Network) | Neurology | 6;81(6):559-565 | 2013 |

| | | | | | |
|-----|---|--|--|-----------------------------------|------|
| 432 | Sokal R, Fleming KM & Tata LJ | Potential of general practice data for congenital anomaly research: Comparison with registry data in the united kingdom | Birth Defects Res A Clin Mol Teratol | 97(8):546-53 | 2013 |
| 431 | Edwards IR, Karimi G, Bergvall T, Caster O, Asiimwe A, Star K, Bhayat F, Soriano-Gabarro M, Thompson M, Bate A & Norén GN | Risk identification in Healthcare Records: comparison to epidemiological studies | 29th International Conference on Pharmacoepidemiology and Therapeutic Risk Management (ICPE) | Montreal, Canada | 2013 |
| 430 | Liu K, Hauben M & Bate A | Signal Refinement of Drug-Induced Thrombocytopenia using The Health Information Network (THIN) Database | 29th International Conference on Pharmacoepidemiology and Therapeutic Risk Management (ICPE) | Montreal, Canada | 2013 |
| 429 | Khalid JM, Globe G, Fox KM, Chau D, Maguire A & Chiou CF | Treatment and referral patterns for psoriasis in United Kingdom primary care: a retrospective cohort study | BMC Dermatol | doi: 10.1186/1471-5945-13-9 | 2013 |
| 428 | Mook P, Johnson M, Thompson M & Bourke A | Comparison of general practices (and their patients) that contribute additional information to The Health Improvement Network (THIN) primary care dataset regularly and those who do so on an ad hoc basis | Society for Academic Primary Care (SAPC) | Nottingham, UK | 2013 |
| 427 | Martín Merino E, Johansson S, Nagy P & García Rodríguez LA | Effect of Baseline Gastrointestinal Risk and Use of Proton Pump Inhibitors on Frequency of Discontinuation of Aspirin for Secondary Cardiovascular Prevention in United Kingdom Primary Care | American Journal of Cardiology | S0002-9149(13)01276-9 | 2013 |
| 426 | Man A, Zhu Y, Zhang Y, Dubreuil M, Rho YH, Peloquin C, Simms RW & Choi HK | The risk of cardiovascular disease in systemic sclerosis: a population-based cohort study | Annals of the Rheumatic Diseases | 72(7):1188-93 | 2013 |
| 425 | Garcia Rodriguez LA, Johansson S & Cea Soriano L | Use of clopidogrel and proton pump inhibitors after a serious acute coronary event: Risk of coronary events and peptic ulcer bleeding | Journal of Thrombosis and Haemostasis | 110(5):1014-24 | 2013 |
| 424 | James GD, Petersen I, Nazareth I, Wedzicha JA & Donaldson GC | Use of long-term antibiotic treatment in COPD patients in the UK: a retrospective cohort study | Primary Care Respiratory Journal | doi: 10.4104/pcrj.2013.00061 | 2013 |
| 423 | List R, Sedani T, Johnson M & Lucrezi G | What impact has NICE clinical guideline 69 had on the diagnosis of, and antibiotic prescribing for, upper respiratory tract infections? A THIN database study | Society for Academic Primary Care (SAPC) | Nottingham, UK | 2013 |
| 422 | Huang Y, Britton J, Hubbard R, Lewis S | Who receives prescriptions for smoking cessation medications? An association rule mining analysis using a large primary care database | Tob Control | 22(4):274-9 | 2013 |
| 421 | Cea-Soriano L, Johansson S & García Rodríguez LA | Risk factors for falls with use of acid-suppressive drugs | Epidemiology | doi: 10.1097/EDE.0b013e318294bec6 | 2013 |
| 420 | Douglas L & Szatkowski L | Socioeconomic variations in access to smoking cessation interventions in UK primary care: insights using the Mosaic classification in a large dataset of primary care records | BMC Public Health | 5;13:546 | 2013 |
| 419 | Shah SM, Carey IM, Harris T, Dewilde S, Victor CR & Cook DG | The effect of unexpected bereavement on mortality in older couples | American Journal of Public Health | 103(6):1140-5 | 2013 |
| 418 | Carey IM, Shah SM, Harris T, Dewilde S & Cook DG | A new simple primary care morbidity score predicted mortality and better explains between practice variations than the Charlson index | Journal of Clinical Epidemiology | 66(4):436-444 | 2013 |

| | | | | | |
|-----|---|---|--|------------------------|------|
| 417 | Hakimi Z, Johnson M, Thompson M, Nazir J, Blak BT & Odeyemi I | Drug treatment patterns for the management of men with benign prostatic hyperplasia (BPH) who have predominant urgency and frequency symptoms in UK primary care The Health Improvement Network (THIN) data | 18th Annual International Meeting (ISPOR) | New Orleans, USA | 2013 |
| 416 | Hakimi Z, Johnson M, Thompson M, Nazir J, Blak BT & Odeyemi IA | Drug treatment patterns for the management of men with benign prostatic hyperplasia (BPH) who have predominant urgency and frequency symptoms in uk primary care the health improvement network (THIN) data. | Value in Health | 16(3):A184 | 2013 |
| 415 | Horsfall LJ, Nazareth I, Pereira SP & Petersen I | Gilbert's syndrome and the risk of death: a population-based cohort study. | Journal of Gastroenterology and Hepatology | doi: 10.1111/jgh.12279 | 2013 |
| 414 | Shah SM, Carey IM, Harris T, Dewilde S & Cook DG | Mortality in older care home residents in England and Wales | Age and Ageing | 42(2):209-215 | 2013 |
| 413 | Murray ML, Hsia Y, Glaser K, Simonoff E, Murphy DG, Asherson PJ, Eklund H & Wong IC | Pharmacological treatments prescribed to people with autism spectrum disorder (ASD) in primary health care | Psychopharmacology | [ahead of print] | 2013 |
| 412 | Vial D, Thompson M, Cockbain C, Hogan S, Johnson M & Bourke A | Rapid identification and recruitment of patients from The Health Improvement Network (THIN) primary care patient data for a health-related quality of life (HRQoL) study of patients with atrial fibrillation (AF) in the UK | 18th Annual International Meeting (ISPOR) | New Orleans, USA | 2013 |
| 411 | Vial D, Thompson M, Cockbain C, Hogan S, Johnson M & Bourke A | Rapid identification and recruitment of patients from the health improvement network (thin) primary care patient data for a health-related quality of life (HRQOL) study of patients with atrial fibrillation (AF) in the United Kingdom. | Value in Health | 16(3):A45-6 | 2013 |
| 410 | Musah A, Gibson JE, Leonardi-Bee J, Cave MR, Ander EL & Bath-Hextall F | Regional variations of Basal cell carcinoma incidence in the UK using THIN database (2004-2010) | British Journal of Dermatology | doi: 10.1111/bjd.12446 | 2013 |
| 409 | Dhalwani NN, Fiaschi L, West J, Tata LJ | Occurrence of fertility problems presenting to primary care: population-level estimates of clinical burden and socioeconomic inequalities across the UK | Hum Reprod | 28(4):960-8 | 2013 |
| 408 | Fardet L, Nazareth I, Whitaker HJ & Petersen I | Severe neuropsychiatric outcomes following discontinuation of long-term glucocorticoid therapy: a cohort study | Journal of Clinical Psychiatry | 74(4):e281-6 | 2013 |
| 407 | Murray ML, Insuk S, Banaschewski T, Neubert AC, McCarthy S, Buitelaar JK, Coghill D, Dittmann RW, Konrad K, Panei P, Rosenthal E, Sonuga-Barke EJ & Wong IC | An inventory of European data sources for the long-term safety evaluation of methylphenidate | European Child & Adolescent Psychiatry | 22(10):605-18 | 2013 |
| 406 | Gao S, Juhaeri J, Reshef S & Dai WS | Association between body mass index and suicide, and suicide attempt among british adults: The health improvement network database | Obesity | 21(3):E334-42 | 2013 |
| 405 | Meeraus WH, Petersen I, Chin RF, Knott F, Gilbert R | Childhood epilepsy recorded in primary care in the UK | Arch Dis Child | 98(3):195-202 | 2013 |
| 404 | Panca M, Cutting K & Guest JF | Clinical and cost-effectiveness of absorbent dressings in the treatment of highly exuding VLUs | Journal of Wound Care | 22(3):109-10, 112-8 | 2013 |
| 403 | Collins GS & Altman DG | Identifying patients with undetected renal tract cancer in primary care: An independent and external validation of Qcancer(®) (Renal) prediction model | Cancer Epidemiology | 37(2):115-20 | 2013 |

| | | | | | |
|-----|--|---|---|---------------------------------------|------|
| 402 | Petherick ES, Cullum NA & Pickett KE | Investigation of the effect of deprivation on the burden and management of venous leg ulcers: a cohort study using the THIN database | Public Library of Science (PLoS one) | 8(3):e58948 | 2013 |
| 401 | Ogdie A, Langan S, Love T, Haynes K, Shin D, Seminara N, Mehta NN, Troxel A, Choi H & Gelfand JM | Prevalence and treatment patterns of psoriatic arthritis in the UK | Rheumatology (Oxford) | 52(3):568-75 | 2013 |
| 400 | Szatkowski L & McNeill A | The delivery of smoking cessation interventions to primary care patients with mental health problems | Addiction | 108(8):1487-94 | 2013 |
| 399 | Zhou X, Murugesan S, Bhullar H, Liu Q, Cai B, Wentworth C & Bate A | An Evaluation of the THIN Database in the OMOP Common Data Model for Active Drug Safety Surveillance | Drug Safety | 36(2):119-34 | 2013 |
| 398 | McCarthy S, Wilton L, Murray M, Hodgkins P, Asherson P & Wong IC | Management of adult attention deficit hyperactivity disorder in UK primary care: a survey of general practitioners | Health and Quality of Life Outcomes | 22;11(1):22 | 2013 |
| 397 | Obiora E, Hubbard R, Sanders RD, Myles PR | The impact of benzodiazepines on occurrence of pneumonia and mortality from pneumonia: a nested case-control and survival analysis in a population-based cohort | Thorax | 68(2):163-70 | 2013 |
| 396 | Tannen R, Xie D, Wang X, Yu M, Weiner MG | A new "Comparative Effectiveness" assessment strategy using the THIN database: comparison of the cardiac complications of pioglitazone and rosiglitazone. | Pharmacoepidemiol Drug Saf | 22(1):86-97 | 2013 |
| 395 | King M, Vasanthan M, Petersen I, Jones L, Marston L & Nazareth I | Mortality and medical care after bereavement: a general practice cohort study | Public Library of Science (PLoS one) | 8(1):e52561 | 2013 |
| 394 | Hall GC, McMahan AD, Dain MP, Wang E & Home PD | Primary-care observational database study of the efficacy of GLP-1 receptor agonists and insulin in the UK | Diabetic Medicine | 30(6):681-86 | 2013 |
| 393 | Powell HA, Iyen-Omofoman B, Hubbard RB, Baldwin DR & Tata LJ | The association between smoking quantity and lung cancer in men and women | Chest | 143(1):123-9 | 2013 |
| 392 | Hwang A, Narayan V & Yang YX | Type 2 diabetes mellitus and survival in pancreatic adenocarcinoma: a retrospective cohort study | Cancer | 15;119(2):404-10 | 2013 |
| 391 | Man SL, Petersen I, Thompson M & Nazareth I | Antiepileptic drugs during pregnancy in primary care: a UK population based study | Public Library of Science (PLoS one) | 7(12):e52339 | 2012 |
| 390 | Wall H, Smith C, Hubbard R | Body mass index and obstructive sleep apnoea in the UK: a cross-sectional study of the over-50s | Prim Care Resp J | 21(4):371-376 | 2012 |
| 389 | Guest JF, Bai JJ, Taylor RR, Sladkevicius E, Lee JP & Lachmann RH | Costs and outcomes over 36 years of patients with phenylketonuria who do and do not remain on a phenylalanine-restricted diet | Journal of Intellectual Disability Research | doi: 10.1111/j.1365-2788.2012.01568.x | 2012 |
| 388 | Shah SM, Carey IM, Harris T, Dewilde S, Victor CR & Cook DG | Do good health and material circumstances protect older people from the increased risk of death after bereavement? | American Journal of Epidemiology | 176(8):689-698 | 2012 |
| 387 | McCarthy S, Wilton L, Murray ML, Hodgkins P, Asherson P & Wong IC | Persistence of pharmacological treatment into adulthood, in UK primary care, for ADHD patients who started treatment in childhood or adolescence | BMC Psychiatry | 5;12:219 | 2012 |
| 386 | Horsfall L, Walters K & Petersen I | Identifying periods of acceptable computer usage in primary care research databases | Pharmacoepidemiology and Drug Safety | [ahead of print] | 2012 |
| 385 | Raman SR, Marshall SW, Haynes K, Gaynes B, Naftel AJ & Stürmer T | Stimulant treatment and injury among children with Attention Deficit Hyperactivity Disorder: an application of the self-controlled case series study design | Injury Prevention | [ahead of print] | 2012 |
| 384 | Currie CJ, Peyrot M, Morgan CL, Poole CD, Jenkins-Jones S, Rubin RR, Burton CM & Evans M | The impact of treatment non-compliance on mortality in people with type 1 diabetes | Journal of Diabetes and its Complications | S1056-8727(12)00286-3 | 2012 |

| | | | | | |
|-----|--|--|---|---------------------------|------|
| 383 | Choi H K, Rho Y H, Zhu Y, Soriano L C, Avina-Zubieta A & Zhang Y | The Risk of Pulmonary Embolism and Deep Vein Thrombosis in Rheumatoid Arthritis: A UK Population-Based Outpatient Cohort Study | Annals of the Rheumatic Diseases | [ahead of print] | 2012 |
| 382 | Woodman J, Freemantle N, Allister J, de Lusignan S, Gilbert R & Petersen I | Variation in recorded child maltreatment concerns in UK primary care records: a cohort study using The Health Improvement Network (THIN) database | Public Library of Science (PLoS one) | 7(11):e49808 | 2012 |
| 381 | Tannen R, Xie D, Wang X, Yu M & Weiner MG | A new "Comparative Effectiveness" assessment strategy using the THIN database: comparison of the cardiac complications of pioglitazone and rosiglitazone. | Pharmacoepidemiology and Drug Safety | 22(1):86-96 | 2012 |
| 380 | Kronman MP, Zaoutis TE, Haynes K, Feng R & Coffin SE | Antibiotic Exposure and IBD Development Among Children: A Population-Based Cohort Study | Pediatrics | 130(4):e794-803 | 2012 |
| 379 | Horsfall LJ, Nazareth I & Petersen I | Cardiovascular Events as a Function of Serum Bilirubin Levels in a Large Statin-Treated Cohort | Circulation | [ahead of print] | 2012 |
| 378 | Gore M, Brix Finnerup N, Sadosky A, Tai KS, Cappelleri JC, Mardekian J, George Rice C & Nieshoff E | Pain-related pharmacotherapy, healthcare resource use and costs in spinal cord injury patients prescribed pregabalin | Spinal Cord | [ahead of print] | 2012 |
| 377 | Sedani T, Bourke A, Thompson M & Johnson M | Seasonal trends of labyrinthitis in the UK – A THIN database study | Society for Academic Primary Care (SAPC) | Glasgow, Scotland | 2012 |
| 376 | Gutiérrez L, Roskell N, Castellsague J, Beard S, Rycroft C, Abeyasinghe S, Shannon P, Gitlin M & Robbins S | Clinical burden and incremental cost of fractures in postmenopausal women in the United Kingdom | Bone | 51(3):324-31 | 2012 |
| 375 | Wise BL, Peloquin C, Choi H, Lane NE & Zhang Y | Impact of Age, Sex, Obesity, and Steroid Use on Quinolone-associated Tendon Disorders | The American Journal of Medicine | 125(12):1228.e23-1228.e27 | 2012 |
| 374 | Osborn DP, Horsfall L, Hassiotis A, Petersen I, Walters K & Nazareth I | Access to cancer screening in people with learning disabilities in the UK: cohort study in the health improvement network, a primary care research database. | Public Library of Science (PLoS one) | 7(8):e43841 | 2012 |
| 373 | Mamtani R, Haynes K, Bilker WB, Vaughn DJ, Strom BL, Glanz K & Lewis JD | Association Between Longer Therapy With Thiazolidinediones and Risk of Bladder Cancer: A Cohort Study | Journal of the National Cancer Institute | 19;104(18):1411-21 | 2012 |
| 372 | Haynes K, Fortier K, Newcomb C, Roy J, Ming E, Horne L, Wood-Ives J, Strom B & Lo Re V | Evaluation of Methods to Estimate Days' Supply of Oral Antidiabetic Drugs in The Health Improvement Network | 28th International Conference on Pharmacoepidemiology and Therapeutic Risk Management | Barcelona, Spain | 2012 |
| 371 | Jon Love T, Zhu Y, Zhang Y, Wall-Burns L, Ogdie A, Gelfand JM & Choi HK | Obesity and the risk of psoriatic arthritis: a population-based study | Annals of the Rheumatic Diseases | 71(8):1273-7 | 2012 |
| 370 | Guest JF, Taylor RR, Vowden K & Vowden P | Relative cost-effectiveness of a skin protectant in managing venous leg ulcers in the UK | Journal of wound care | 21(8):389-94, 396-7 | 2012 |
| 369 | Rho YH, Zhu Y, Zhang Y, Reginato AM & Choi HK | Risk factors for pseudogout in the general population | Rheumatology (Oxford) | 51(11):2070-4 | 2012 |
| 368 | List R, Sedani T, Agustini S & Lucrezi G | What Impact Has National Institute for Clinical Excellence (NICE) Clinical Guideline 69 Had On The Diagnosis and Antibiotic Prescribing For Upper Respiratory Tract Infections? An Analysis Of The THIN Database | 28th International Conference on Pharmacoepidemiology and Therapeutic Risk Management | Barcelona, Spain | 2012 |
| 367 | Blak BT, Smith HT, Hards M, Maguire A & Gimeno V | A retrospective database study of insulin initiation in patients with Type 2 diabetes in UK primary care | Diabetic Medicine | 29(8): e191-e198 | 2012 |

| | | | | | |
|-----|--|---|---|-----------------------------------|------|
| 366 | Woodman J, Allister J, Rafi I, de Lusignan S, Belsey J, Petersen I & Gilbert R | A simple approach to improve recording of concerns about child maltreatment in primary care records: developing a quality improvement intervention | British Journal of General Practice | 62(600):478-86 | 2012 |
| 365 | Berry SD, Zhu Y, Choi H, Kiel DP & Zhang Y | Diuretic initiation and the acute risk of hip fracture | Osteoporosis International | [ahead of print] | 2012 |
| 364 | Ashman K E , Cawood A L & Stratton R J | OC-038 Healthcare use according to body mass index (BMI) category in individuals registered to GP practices | An International Journal of Gastroenterology and Hepatology - BMJ | 61:A16-A17 | 2012 |
| 363 | Blak BT, Smith HT, Hards M, Curtis BH & Ivanyi T | Optimization of insulin therapy in patients with Type 2 diabetes mellitus: beyond basal insulin | Diabetic Medicine | 29(7):e13-20 | 2012 |
| 362 | Collins GS & Altman DG | Predicting the adverse risk of statin treatment: an independent and external validation of Qstatin risk scores in the UK | Heart | 98(14):1091-7 | 2012 |
| 361 | Fardet L, Petersen I & Nazareth I | Risk of cardiovascular events in people prescribed glucocorticoids with iatrogenic Cushing's syndrome: cohort study | British Medical Journal (BMJ) | 345:e4928. doi: 10.1136/bmj.e4928 | 2012 |
| 360 | Lo Re V 3rd, Haynes K, Ming EE, Wood Ives J, Horne LN, Fortier K, Carbonari DM, Hennessy S, Cardillo S, Reese PP, Reddy KR, Margolis D, Apter A, Kimmel SE, Roy J, Freeman CP, Razzaghi H, Holick CN, Esposito DB, Van Staa TP, Bhullar H & Strom BL | Safety of saxagliptin: rationale for and design of a series of postmarketing observational studies | Pharmacoepidemiology and Drug Safety | 10.1002/pds.3318 | 2012 |
| 359 | Iyen-Omofoman B, Hubbard RB, Baldwin DR & Tata LJ | The association between smoking quantity and lung cancer in men and women | Chest | 10.1378/chest.12-1068. | 2012 |
| 358 | Idris I, Warren G & Donnelly R | Association Between Thiazolidinedione Treatment and Risk of Macular Edema Among Patients With Type 2 DiabetesThiazolidinedione and Macular Edema in Type 2 DM | Archives of Internal Medicine | 11:1-7 | 2012 |
| 357 | Wall H, Smith C & Hubbard R | BMI and obstructive sleep apnoea in the UK: a crosssectional study of the over-50s | Primary Care Respiratory Journal | 10.4104/pcrj.2012.00053 | 2012 |
| 356 | Renehan AG, Yeh HC, Johnson JA, Wild SH, Gale EA, Moller H, Diabetes and Cancer Research Consortium | Diabetes and cancer (2): evaluating the impact of diabetes on mortality in patients with cancer. | Diabetologia | 55(6):1619-32 | 2012 |
| 355 | Johnson JA, Carstensen B, Witte D, Bowker SL, Lipscombe L & Renehan AG, | Diabetes and cancer (1): evaluating the temporal relationship between type 2 diabetes and cancer incidence | Diabetologia | 55(6):1607-18 | 2012 |
| 354 | Ruigómez A, Johansson S, Wernersson B, Fernández Cantero O & García Rodríguez LA | Gastroesophageal reflux disease in primary care: using changes in proton pump inhibitor therapy as an indicator of partial response | Scandinavian Journal of Gastroenterology | 47(7):751-61 | 2012 |
| 353 | Collins GS & Altman DG | Predicting the 10 year risk of cardiovascular disease in the United Kingdom: independent and external validation of an updated version of QRISK2 | British Medical Journal (BMJ) | 21;344:e4181 | 2012 |
| 352 | McCarthy S, Wilton L, Murray ML, Hodgkins P, Asherson P & Wong IC | The epidemiology of pharmacologically treated attention deficit hyperactivity disorder (ADHD) in children, adolescents and adults in UK primary care | BMC Pediatrics | 19;12(1):78 | 2012 |
| 351 | McCarthy S, Wilton L, Murray ML, Hodgkins P, Asherson P & Wong IC | The epidemiology of pharmacologically treated attention deficit hyperactivity disorder (ADHD) in children, adolescents and adults in UK primary care | BMC Pediatrics | 19;12(1):78 | 2012 |
| 350 | Article | UK data link glitazones with diabetic macular oedema | Pharmaceutical Journal | 288:701 | 2012 |

| | | | | | |
|-----|---|--|---|---------------------------------------|------|
| 349 | Jara M, Wentworth C 3rd & Lanes S | A new user cohort study comparing the safety of long-acting inhaled bronchodilators in COPD | BMJ Open | 22;2(3). pii: e000841 | 2012 |
| 348 | Rowan CG, Brunelli SM, Munson J, Flory J, Reese PP, Hennessy S, Lewis J, Mines D, Barrett JS, Bilker W & Strom BL | Clinical importance of the drug interaction between statins and CYP3A4 inhibitors: a retrospective cohort study in The Health Improvement Network | Pharmacoepidemiology and Drug Safety | 21(5):494-506 | 2012 |
| 347 | Belsey JD | Optimizing adherence in hypertension: a comparison of outcomes and costs using single tablet regimens vs individual component regimens | Journal of Medical Economics | doi:10.3111/13696998.2012.689791 | 2012 |
| 346 | Walters K, Rait G, Hardoon S, Kalaitzaki E, Petersen I & Nazareth I | Socio-demographic variation in chest pain incidence and subsequent coronary heart disease in primary care in the United Kingdom | European Journal of Preventive Cardiology | [ahead of print] | 2012 |
| 345 | Fardet L, Petersen I & Nazareth I | Suicidal behavior and severe neuropsychiatric disorders following glucocorticoid therapy in primary care | American Journal of Psychiatry | 169(5):491-7 | 2012 |
| 344 | Langley T E, McNeill A, Lewis S, Szatkowski L & Quinn C | The impact of media campaigns on smoking cessation activity: A structural vector autoregression analysis | Addiction | doi: 10.1111/j.1360-0443.2012.03958.x | 2012 |
| 343 | Taggar JS, Coleman T, Lewis S & Szatkowski L | The impact of the Quality and Outcomes Framework (QOF) on the recording of smoking targets in primary care medical records: Cross-sectional analyses from The Health Improvement Network (THIN) database | BMC Public Health | 12;329 | 2012 |
| 342 | Cai B, Xu W, Bortnichak E & Watson DJ | An algorithm to identify medical practices common to both the General Practice Research Database and The Health Improvement Network database | Pharmacoepidemiology and Drug Safety | 21(7):770-774 | 2012 |
| 341 | Woodman J, et al. | Child maltreatment-related concerns recorded in UK primary care: a study using the THIN database | 16th April 2012 BASCPAN | Belfast | 2012 |
| 340 | Orton E, Kendrick D, West J & Tata LJ | Independent risk factors for injury in pre-school children: three population-based nested case-control studies using routine primary care data. | Public Library of Science (PLoS one) | 2012;7(4):e35193 | 2012 |
| 339 | Flory JH, Ky B, Haynes K, M Brunelli S, Munson J, Rowan C, Strom BL & Hennessy S | Observational cohort study of the safety of digoxin use in women with heart failure | BMJ Open | 13;2(2):e000888 | 2012 |
| 338 | Collins G & Altman D | Predicting the risk of chronic kidney disease in the UK: an evaluation of QKidney [®] scores using a primary care database | British Journal of General Practice | 62(597):e243-50 | 2012 |
| 337 | Forde KA, Haynes K, Troxel AB, Trooskin S, Osterman MT, Kimmel SE, Lewis JD & Lo Re V 3rd | Risk of myocardial infarction associated with chronic hepatitis C virus infection: a population-based cohort study | Journal of viral hepatitis | 19(4):271-7 | 2012 |
| 336 | Martín-Merino E, Fortuny J, Rivero E & García-Rodríguez LA | Validation of diabetic retinopathy and maculopathy diagnoses recorded in a U.K. primary care database | Diabetes Care | 35(4):762-7 | 2012 |
| 335 | GC Hall, McMahan, D Carroll & PD Home | Macrovascular and microvascular outcomes after beginning of insulin versus additional oral glucose-lowering therapy in people with type 2 diabetes: an observational study | Pharmacoepidemiology and Drug Safety | 21(3):305-13 | 2012 |
| 334 | Wijlaars LP, Nazareth I & Petersen I | Trends in depression and antidepressant prescribing in children and adolescents: a cohort study in The Health Improvement Network (THIN). | Public Library of Science (PLoS one) | 7(3):e33181 | 2012 |
| 333 | Meropol SB & Metlay JP | Accuracy of pneumonia hospital admissions in a primary care electronic medical record database | Pharmacoepidemiology and Drug Safety | doi: 10.1002/pds.3207 | 2012 |

| | | | | | |
|-----|---|--|---|---------------------------------------|------|
| 332 | Taylor RR, Sladkevicius E, Panca M, Lack G & Guest JF | Cost-effectiveness of using an extensively hydrolysed formula compared to an amino acid formula as first-line treatment for cow milk allergy in the UK | Pediatric Allergy and Immunology | 23(3):240-9 | 2012 |
| 331 | Trifirò G, Mokhles MM, Dieleman JP, van Soest EM, Verhamme K, Mazzaglia G, Herings R, de Luise C, Ross D, Brusselle G, Colao A, Haverkamp W, Schade R, van Camp G, Zanettini R & Sturkenboom MC | Risk of cardiac valve regurgitation with dopamine agonist use in Parkinson's disease and hyperprolactinaemia: a multi-country, nested case-control study | Drug Safety | 35(2):159-71 | 2012 |
| 330 | F Lapi, F. Nicotra, L. Scotti, A. Vannacci, M Thompson, F Pieri, N Mugelli, A Zambon, G Corrao, A Mugelli & A Rubino | Use Of Antidepressant Serotonergic Medications And Cardiac Valvulopathy: A Nested Case-Control Study In The Health Improvement Network (Thin) Database | British Journal of Clinical Pharmacology | 74(3):536-544 | 2012 |
| 329 | Lapi F, Nicotra F, Scotti L, Vannacci A, Thompson M, Pieri F, Mugelli N, Zambon A, Corrao G, Mugelli A & Rubino A | Use of antidepressants serotonergic medications and cardiac valvulopathy: A nested case-control study in The Health Improvement Network (THIN) database | British Journal of Clinical Pharmacology | doi: 10.1111/j.1365-2125.2012.04224.x | 2012 |
| 328 | L Shallcross | A hidden epidemic of impetigo – trends from the THIN database of primary care consultations | Infectious Disease Research Network | | 2012 |
| 327 | Harris T, Shah SM, Carey IM, DeWilde S & Cook DG | Antidepressant prescribing in older primary care patients in community and care home settings in England and Wales | Journal of the American Medical Directors Association | 13:41-7 | 2012 |
| 326 | Choi HK, Soriano LC, Zhang Y & Rodríguez LA | Calcium Channel Blockers, Losartan, and the Risk of Incident Gout among Patients with Hypertension: A Population-Based Case-Control Study | British Medical Journal (BMJ) | 344:d8190. doi: 10.1136/bmj.d8190 | 2012 |
| 325 | Man S, Petersen I, Thompson M & Nazareth I | Discontinuation of antiepileptic drugs in pregnancy; a study in The Health Improvement Network (THIN) primary care database | Infectious Disease Research Network | | 2012 |
| 324 | Hall GC, McMahon AD, Dain M-P & et al | GLP-1 Receptor Agonist (GLP1ra) and Insulin Efficacy in Primary Health Care | Diabetes | 61(S1):A276 | 2012 |
| 323 | Menon S, Dhar A, Hoare J & Trudgill N | How commonly is gastric cancer missed at endoscopy (a UK primary care based study) ? | Digestive Disease Week conference | San Diego, USA | 2012 |
| 322 | Hall GC, McMahon AD, Carroll D & et al | Observational Study of the Association of First Insulin Type in Uncontrolled Type 2 Diabetes with Macrovascular and Microvascular Disease | Public Library of Science (PLoS one) | 7(11):e49908 | 2012 |
| 321 | Langley TE, Szatkowski L, McNeill A, Coleman T & Lewis S | Prescribing of nicotine replacement therapy to cardiovascular disease patients in England | Addiction | 107(7):1341-8 | 2012 |
| 320 | Shah SM, Carey IM, Harris T, DeWilde S & Cook DG | Quality of Prescribing in Care Homes and the Community in England and Wales | British Journal of General Practice | 62, e329-e336 | 2012 |
| 319 | Walters K, Rait G, Griffin M, Buszewicz M & Nazareth I | Recent trends in the incidence of anxiety diagnoses and symptoms in primary care | Public Library of Science (PLoS one) | 7(8):e41670 | 2012 |
| 318 | Shah SM, Carey IM, Harris T, Dewilde S & Cook DG | The impact of dementia on influenza vaccination uptake in community and care home residents | Age and Ageing | 41(1):64-9 | 2012 |
| 317 | P L Thompson P L, Chretien S, Bugnard F, Maier W C, Ito T, Spencer M & Thompson G | The use of GP questionnaires to assess psoriasis severity in The Health Improvement Network (THIN) database | Infectious Disease Research Network | | 2012 |
| 316 | Gore M, Sadosky AB, Leslie DL, Tai KS & Emery P | Therapy Switching, Augmentation, and Discontinuation in Patients with Osteoarthritis and Chronic Low Back Pain | Pain Practice | 12(6):457-68 | 2012 |

| | | | | | |
|-----|---|---|--|---|------|
| 315 | Wijlaars L, Irwin Nazareth I & Irene Petersen I | Trends in antidepressant prescribing in children in the United Kingdom, a study in The Health Improvement Network (THIN) | Infectious Disease Research Network | | 2012 |
| 314 | Thompson M, Dattani H, Bourke A, MacLachlan S, Lloyd D & Faulding S | A comparison between prescribing in THIN data and dispensing in England | EuroDURG/ISPE meeting | University of Antwerp | 2011 |
| 313 | Man SL | Better public health through pharmaco-epidemiology and quality use of medicine | EuroDURG/ISPE meeting | University of Antwerp | 2011 |
| 312 | Blak BT, Thompson M, Dattani H & Bourke A | Generalisability of The Health Improvement Network (THIN) database: demographics, chronic disease prevalence and mortality rates. | Informatics in Primary Care | 19(4):251-5 | 2011 |
| 311 | Hayes J, Prah P, Nazareth I, King M, Walters K, Petersen I & Osborn D | Prescribing trends in bipolar disorder: cohort study in the United Kingdom THIN primary care database 1995-2009 | Public Library of Science (PLoS one) | 2011;6(12):e28725 | 2011 |
| 310 | White D, Choi H, Peloquin C, Zhu Y & Zhang Y | Secular trend of adhesive capsulitis | Arthritis Care & Research | 63(11):1571-5 | 2011 |
| 309 | Gribbin J, Hubbard R, Gladman J, Smith C & Lewis S | Serotonin-norepinephrine reuptake inhibitor antidepressants and the risk of falls in older people: case-control and case-series analysis of a large UK primary care database. | Drugs Aging | 28(11):895-902 | 2011 |
| 308 | Iyem-Omofoman B, Hubbard RB, Smith CJ, Sparks E, Bradley E, Bourke A & Tata LJ | The distribution of lung cancer across sectors of society in the United Kingdom: a study using national primary care data | BMC Public Health | 10;11:857 | 2011 |
| 307 | Blak BT, Lee J & Dungarwalla M | Trends in UK smoking cessation prescription expenditure over time - a THIN database study | 14th Annual European Congress (ISPOR) | Madrid, SPAIN | 2011 |
| 306 | Denburg MR, Haynes K, Shults J, Lewis JD & Leonard MB | Validation of The Health Improvement Network (THIN) database for epidemiologic studies of chronic kidney disease | Pharmacoepidemiology and Drug Safety | 20(11):1138-49. doi: 10.1002/pds.2203 | 2011 |
| 305 | Thompson M, Hards M & Blak BT | Waist circumference and body mass index recording – a THIN database study | 14th Annual European Congress (ISPOR) | Madrid, SPAIN | 2011 |
| 304 | Langley TE, Szatkowski LC, Wythe S & Lewis SA | Can primary care data be used to monitor regional smoking prevalence? An analysis of The Health Improvement Network primary care data | BMC Public Health | 7;11:773 | 2011 |
| 303 | Gribbin J, Hubbard R, Gladman J, Smith C & Lewis S | Serotonin-Norepinephrine Reuptake Inhibitor Antidepressants and the Risk of Falls in Older People: Case-Control and Case-Series Analysis of a Large UK Primary Care Database | Drugs Aging | 28(11):895-902 | 2011 |
| 302 | Loftus J, Christopoulou D, Blak BT & Hards M | Topical prostaglandin fixed combinations in UK primary care: observational study using data from the health improvement network | European Journal of Ophthalmology | doi: 10.5301/ejo.5000068 | 2011 |
| 301 | Navaratnam V, Ali N, Smith CJ, McKeever T, Fogarty A & Hubbard RB | Does the presence of connective tissue disease modify survival in patients with pulmonary fibrosis? | Respiratory Medicine | 105(12):1725-1930 | 2011 |
| 300 | David Lloyd - The NHS Information Centre, Prescribing and Primary Care Services | Prescribing compliance: a review of the proportion of prescriptions dispensed | The Health and Social Care Information Centre | Report [www.ic.nhs.uk/pubs/presccompliance] | 2011 |
| 299 | Thompson M, Dattani H, Bourke A, MacLachlan S, Lloyd D & Faulding S | A comparison between prescribing in THIN data and dispensing in England | 27th International conference on pharmacoepidemiology & therapeutic risk management (ICPE) | Chicago, USA | 2011 |
| 298 | García Rodríguez LA, Cea-Soriano L, Martín-Merino E & Johansson S | Discontinuation of low dose aspirin and risk of myocardial infarction: case-control study in UK primary care. | British Medical Journal (BMJ) | 343:d4094. doi: 10.1136/bmj.d4094 | 2011 |
| 297 | Douglas I, Evans S & Smeeth L | Effect of statin treatment on short term mortality after pneumonia episode: cohort study | British Medical Journal (BMJ) | 342:d1642. doi: 10.1136/bmj.d1642. | 2011 |

| | | | | | |
|-----|--|--|--|-----------------------|------|
| 296 | Marshall T, Dattani H, Thompson M, Bourke A, Pinder L & Blak BT | Linking cancer registry records to THIN - a UK general practice database | 27th International conference on pharmacoepidemiology & therapeutic risk management (ICPE) | Chicago, USA | 2011 |
| 295 | Prah P, Petersen I, Nazareth I, Walters K & Osborn D | National changes in oral antipsychotic treatment for people with schizophrenia in primary care between 1998 and 2007 in the United Kingdom | Pharmacoepidemiology and Drug Safety | doi: 10.1002/pds.2213 | 2011 |
| 294 | Kirby MG, Schnetzler G, Zou KH & Symonds T | Prevalence and detection rate of underlying disease in men with erectile dysfunction receiving phosphodiesterase type 5 inhibitors in the United Kingdom: a retrospective database study | International Journal of Clinical Practice | 65(7):797-806 | 2011 |
| 293 | Douglas I, Evans S & Smeeth L | A cohort study to measure the impact of statin therapy on short-term mortality following a pneumonia episode | Society for Academic Primary Care (SAPC) | Bristol | 2011 |
| 292 | Woodman J, Gilbert R, Allister J, De Lusignan S, Rafi I & Petersen I | Child maltreatment recording in primary care | Society for Academic Primary Care (SAPC) | Bristol | 2011 |
| 291 | Rabi R, Elliott R & Quinn C | Estimating the impact of adherence to allopurinol therapy on the outcomes of gout using The Health Improvement Network (THIN) general practice database | Society for Academic Primary Care (SAPC) | Bristol | 2011 |
| 290 | Orton E, West J, Kendrick & Tata L | Identification pre-school children at risk of poisoning injury using routinely recorded primary care data | Society for Academic Primary Care (SAPC) | Bristol | 2011 |
| 289 | Man S, Nazareth I, Thompson M & Petersen I | Patterns of prescription of antiepileptic drugs in pregnancy, a study in THIN | Society for Academic Primary Care (SAPC) | Bristol | 2011 |
| 288 | Petersen I, Gilbert R, Evans S & Nazareth I | Pregnancy and prescribed psychotropic medicine | Society for Academic Primary Care (SAPC) | Bristol | 2011 |
| 287 | Langley TE, Huang Y, McNeill A, Coleman T, Szatkowski L, Lewis S | Prescribing of smoking cessation medication in England since the introduction of varenicline | Addiction | (7):1319-24 | 2011 |
| 286 | Meeraus W, Petersen I & Gilbert R | Primary -care databases: An alternative to disease registers for population-based research into cerebral palsy in the UK | Society for Academic Primary Care (SAPC) | Bristol | 2011 |
| 285 | Gribbin J, Hubbard R, Gladman J, Smith C & Lewis S | Risk of falls associated with antihypertensive medication: self-controlled case series | Pharmacoepidemiology and Drug Safety | 20(8):879-84 | 2011 |
| 284 | Haynes K, Bilker WB, Tenhave TR, Strom BL & Lewis JD | Temporal and within practice variability in the health improvement network | Pharmacoepidemiology and Drug Safety | 20(9):948-55 | 2011 |
| 283 | Trotter C, Petersen I, Sterne J & Hayward A | The impact of pneumococcal conjugate vaccination on primary care consultations in the UK | Society for Academic Primary Care (SAPC) | Bristol | 2011 |
| 282 | Szatkowski L, Coleman T, McNeill A & Lewis S | The impact of the introduction of smoke-free legislation on prescribing of stop-smoking medications in England | Addiction | 106(10):1827-34 | 2011 |
| 281 | Wijlaars L, Nazareth I & Petersen I | Trends in the incidence of childhood depression in the UK, a study in THIN | Society for Academic Primary Care (SAPC) | Bristol | 2011 |
| 280 | Toh S, García Rodríguez LA & Hernán MA | Confounding adjustment via a semi-automated high-dimensional propensity score algorithm: an application to electronic medical records | Pharmacoepidemiology and Drug Safety | 20(8):849-57 | 2011 |
| 279 | Lassa S, Campbell MJ & Bennett CE | Epidemiology of scabies prevalence in the U.K. from general practice records. | British Journal of Dermatology | 164(6):1329-34 | 2011 |
| 278 | Szatkowski L, McNeill A, Lewis S & Coleman T | A comparison of patient recall of smoking cessation advice with advice recorded in electronic medical records | BMC Public Health | 10;11:291 | 2011 |
| 277 | Szatkowski L, Lewis S, McNeill A, Huang Y & Coleman T | Can data from primary care medical records be used to monitor national smoking prevalence? | Journal of Epidemiology & Community Health | 66(9):791-5. | 2011 |

| | | | | | |
|-----|--|--|--|--------------------|------|
| 276 | Thompson M, Sedani T, Blak BT & Hards M | Impact of free prescriptions in Wales on prescribing - a THIN database study | ISPOR 16th Annual International Meeting | Baltimore, USA | 2011 |
| 275 | Blak BT, Lee J, Thompson M, Bourke A & Pinder L | Is it feasible to collect quality of life data electronically in THIN - a UK general practice database? | ISPOR 16th Annual International Meeting | Baltimore, USA | 2011 |
| 274 | Fardet L, Petersen I & Nazareth I | Description of oral glucocorticoid prescriptions in general population | La Revue de Médecine Interne | 32(10):594-9 | 2011 |
| 273 | Guest JF, Panca M, Baeyens JP, de Man F, Ljungqvist O, Pichard C, Wait S & Wilson L | Health economic impact of managing patients following a community-based diagnosis of malnutrition in the UK | Clinical Nutrition | (4):422-9 | 2011 |
| 272 | de la Iglesia B, Potter JF, Poulter NR, Robins MM & Skinner J | Performance of the ASSIGN cardiovascular disease risk score on a UK cohort of patients from general practice | Heart | 97(6):491-9 | 2011 |
| 271 | Petersen I, Gilbert RE, Evans SJ, Man SL & Nazareth I | Pregnancy as a major determinant for discontinuation of antidepressants: an analysis of data from The Health Improvement Network. | Journal of Clinical Psychiatry | 72(7):979-85 | 2011 |
| 270 | Fardet L, Petersen I & Nazareth I | Prevalence of long-term oral glucocorticoid prescriptions in UK over the past 20 years. | Rheumatology (Oxford) | 50(11):1982-90 | 2011 |
| 269 | Shah SM, Carey IM, Harris T, DeWilde S & Cook DG | Quality of Chronic Disease Care for Older People in Care Homes and the Community in a Primary Care Pay for Performance System: A Retrospective Study | British Medical Journal (BMJ) | BMJ.2011 342: d912 | 2011 |
| 268 | Seminara NM, Abuabara K, Shin DB, Langan SM, Kimmel SE, Margolis D, Troxel AB & Gelfand JM | Validity of The Health Improvement Network (THIN) for the study of psoriasis | British Journal of Dermatology | 164(3):602-9 | 2011 |
| 267 | García Rodríguez L, Lin K J, Sonia Hernández-Díaz S & Johansson S | Risk of Upper Gastrointestinal Bleeding With Low-Dose Acetylsalicylic Acid Alone and in Combination With Clopidogrel and Other Medications | Circulation | 123: 1047-1048 | 2011 |
| 266 | Horsfall L, Rait G, Walters K, Swallow D, Perreira S, Nazareth I & Petersen I | Serum Bilirubin and Risk of Respiratory Disease and Death | Journal of the American Medical Association (JAMA) | 305(7): 691-697 | 2011 |
| 265 | Hall GC, McMahon AD, Dain MP & Home PD | A comparison of duration of first prescribed insulin therapy in uncontrolled type 2 diabetes | Diabetes Research and Clinical Practice | 94(3):442-48 | 2011 |
| 264 | Shah SM, Carey IM, Harris T, DeWilde S & Cook DG | Antipsychotic prescribing to older people living in care homes and the community in England and Wales | International Journal of Geriatric Psychiatry | 26:423-434 | 2011 |
| 263 | Hall GC, Smith HT, Curtis B & McMahon AD | Changes and predictors for change to thiazolidinedione prescribing in UK primary care following the rosiglitazone safety warning | International Journal of Clinical Practice | 65(5):586-91 | 2011 |
| 262 | Harris T, Shah SM, Carey IM, DeWilde S & Cook DG | Depression indicators in a national sample of older community and care home patients: applying the Quality and Outcomes Framework | British Journal of General Practice | 61:135-8 | 2011 |
| 261 | Serumaga B, Ross-Degnan D, Avery AJ, Elliott RA, Majumdar SR, Zhang F & Soumerai SB | Effect of pay for performance on the management and outcomes of hypertension in the United Kingdom: interrupted time series study | British Medical Journal (BMJ) | 25;342:d108 | 2011 |
| 260 | A. Pinçon, G Carrigan, F Sillan, A Khromava & C Pierfitte | From Signal Detection to Evaluation to Action: The experience of Pediaxel® and Apnea | ISOP (International Society on Pharmacovigilance) | Istanbul, Turkey | 2011 |
| 259 | Gutierrez L, Roskell N, Castellsague J, Beard S, Rycroft C, Abesinghe S, Shannon P, Robbins S & Gitlin M | Study of the incremental cost and clinical burden of hip fractures in postmenopausal women in the United Kingdom | Journal of Medical Economics | 14(1): 99-107 | 2011 |
| 258 | Rodríguez LA, Soriano LC & Choi HK | Impact of diabetes against the future risk of developing gout | Annals of the Rheumatic Diseases | 69(12):2090-4 | 2010 |

| | | | | | |
|-----|---|--|---|---------------------------------------|------|
| 257 | Gordon J, Pockett RD, Tetlow AP, McEwan P & Home PD | A comparison of intermediate and long-acting insulins in people with type 2 diabetes starting insulin: an observational database study. | International Journal of Clinical Practice | 64(12):1609-18 | 2010 |
| 256 | Gunathilake W, Song S, Sridharan S, Fernando DJ & Idris I | Cardiovascular and metabolic risk profiles in young and old patients with type 2 diabetes | QJM: An International Journal of Medicine | 103(11):881-4 | 2010 |
| 255 | McBride D, Haroon S, Walters K, Gilmour S & Raine R | Explaining variation in referral from primary to secondary care: cohort study | British Medical Journal (BMJ) | BMJ 2010; 341:c6267 | 2010 |
| 254 | Arana A, Wentworth CE, Fernández-Vidaurre C, Schlienger RG, Conde E & Arellano F | Incidence of cancer in the general population and in patients with or without atopic dermatitis in the U.K | British Journal of Dermatology | 163(5):1036-43 | 2010 |
| 253 | Davé S, Petersen I, Sherr L & Nazareth I | Incidence of maternal and paternal depression in primary care: a cohort study using a primary care database. | Archives of Pediatrics & Adolescent Medicine | 164(11):1038-44 | 2010 |
| 252 | Damery S, Ryan R, Wilson S, Ismail T & Hobbs FR | Iron deficiency anaemia and delayed diagnosis of colorectal cancer: a retrospective cohort study | Colorectal Disease | doi: 10.1111/j.1463-1318.2010.02488.x | 2010 |
| 251 | Szatkowski L, Lewis S, McNeill A & Coleman T | Is smoking status routinely recorded when patients register with a new GP? | Family Practice | 27(6):673-5 | 2010 |
| 250 | Iglesia B, Potter J, Poulter N, Robins M & Skinner J | Performance of the ASSIGN cardiovascular disease risk score on a UK cohort of patients from general practice | Heart BMJ | [ahead of print] | 2010 |
| 249 | Martín-Merino E, Ruigómez A, Johansson S, Wallander MA & García-Rodríguez LA | Study of a cohort of patients newly diagnosed with depression in general practice: prevalence, incidence, comorbidity, and treatment patterns. | Primary Care Companion - Journal of Clinical Psychiatry | 12(1):PCC.08m00764. | 2010 |
| 248 | Kaplan GG, Hubbard J, Korzenik J, Sands BE, Panaccione R, Ghosh S, Wheeler AJ & Villeneuve PJ | The inflammatory bowel diseases and ambient air pollution: a novel association. | The American Journal of Gastroenterology | 105(11):2412-9 | 2010 |
| 247 | Langan SM, Groves RW & West J | The Relationship between Neurological Disease and Bullous Pemphigoid: A Population-Based Case-Control Study. | Journal of Investigative Dermatology | [ahead of print] | 2010 |
| 246 | Petersen I, Gilbert R, Evans S, Ridolfi A & Nazareth I | Oral antibiotic prescribing during pregnancy in primary care: UK population-based study. | Journal of Antimicrobial Chemotherapy | 65(10):2238-46 | 2010 |
| 245 | Owen V, Seetho I & Idris I | Predictors of responders to insulin therapy at 1 year among adults with type 2 diabetes | Diabetes, Obesity & Metabolism | 12(10):865-70 | 2010 |
| 244 | Ralston SH, Kou TD, Wick-Urban B, Steinbuch M & Masud T | Risk of upper gastrointestinal tract events in risedronate users switched to alendronate. | Calcified Tissue International | 87(4):298-304 | 2010 |
| 243 | Gonzalez-Perez A, Schlienger RG & García Rodríguez LA. | Acute pancreatitis in association with type 2 diabetes and anti-diabetic drugs: a population-based cohort study. | Diabetes Care | 33(12):2580-5 | 2010 |
| 242 | Yuan Z, Weinstein R, Zhang J, Cheng M, Griffin G, Zolynas R, Plotnikov AN, Lee MS, Oppenheimer L & Burton P | Antithrombotic therapies in patients with heart failure: hypothesis formulation from a research database. | Pharmacoepidemiology and Drug Safety | 19(9):911-20 | 2010 |
| 241 | Shah SM, Carey IM, Harris T, DeWilde S, Hubbard R, Lewis S & Cook DG | Identifying the clinical characteristics of older people living in care homes using a novel approach in a primary care database. | Age and Ageing | 39(5):617-23 | 2010 |
| 240 | Chacón García A, Ruigómez A & García Rodríguez LA | Incidence rate of community acquired pneumonia in a population cohort registered in BIFAP | Aten Primaria | 42(11):543-9 | 2010 |
| 239 | Feary J R, Rodrigues L, Smith C J, Hubbard R B & Gibson J E | Prevalence of major comorbidities in subjects with COPD and incidence of myocardial infarction and stroke: a comprehensive analysis using data from primary care | Thorax BMJ | 65:956-962 | 2010 |
| 238 | Loftus J, Christopoulou D, Blak BT & Hards M | Use of Topical Prostaglandin Fixed Combination in the UK Primary Care: An Observational Study in The Health Improvement Network (THIN) Database | Congress of the European Glaucoma Society | Madrid, Spain | 2010 |

| | | | | | |
|-----|---|---|--|----------------|------|
| 237 | Blak BT, Gabriel ZL, Hards M & Minhas I | Variations in lipid levels across the UK – a THIN database study | PCCS 13th Annual Scientific Meeting | Leeds, UK | 2010 |
| 236 | Douglas I, Smeeth L & Evans S | A Propensity Score Adjusted Cohort Study To Measure the Impact of Statin Therapy on Short-Term Mortality Following a Pneumonia Episode | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 235 | Meropol S, Localio AR & Metlay JP | Adverse Events Associated with Antibiotic Use | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 234 | Arana A, Arellanom F, Wentworth C & Price J | An Algorithm for the Diagnosis of Suicide in the THIN Database | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 233 | Cai B, Xu W, Bortnichak E & Watson DJ | An algorithm to identify practices common to both the GPRD and THIN Databases. | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 232 | Horsfall L, Rait G, Walters K, Swallow D, Pereira S & Petersen I | Association between Bilirubin and Oxidative Stress Mediated Disease | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 231 | Haynes K, Bilker WB, TenHave TR & Lewis JD | Between and within practice variability among practices within the health improvement network | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 230 | Desamericq G, Laforest L, Schwalm MS, Bourke A, Broquet M, Van Ganse E & Moulin P | Characteristics and LDL-C Therapeutic Objectives of Patients Receiving Unusual Lipid-Modifying Therapy Regimens | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 229 | Blak BT, Wilson K, Metcalfe M, Maguire A, Hards M & Gimeno V | Characteristics of patients receiving varenicline treatment for smoking cessation in UK primary care – a THIN database study | 26th International Conference on Pharmacoepidemiology & Therapeutic Risk Management (ICPE) | Brighton, UK | 2010 |
| 228 | Blak B T, Wilson K, Metcalfe M, Hards M, Maguire A & Gimeno V | Characteristics of Patients Receiving Varenicline Treatment for Smoking Cessation in UK Primary Care–A THIN Database Study | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 227 | Laforest L, Schwalm MS, Dufour M, Broquet M, Bourke A & Van Ganse E | Comparison of Prescribing Patterns for Asthma in Five European Countries | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 226 | Laforest L, Schwalm MS, Dufour M, Bourke A, Broquet M & Van Ganse E | Comparison of prescribing patterns in asthma patients supervised in primary care between five European countries | 26th International Conference on Pharmacoepidemiology & Therapeutic Risk Management (ICPE) | Brighton, UK | 2010 |
| 225 | Prah P, Petersen I, Nazareth I, Walters K & Osborn D | Descriptive analysis of antipsychotic prescriptions in primary care | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 224 | Ruigómez A, Lundborg P, Johansson S, Wallander MA & García Rodríguez LA | Follow-up of a cohort of children and adolescents with gastro-esophageal reflux disease who were free of reflux esophagitis at initial diagnosis. | Scandinavian Journal of Gastroenterology | 45(7-8):814-21 | 2010 |
| 223 | Rothenbacher D, Primatesta P, Ferreira A, Soriano L & Garcia Rodriguez L | Frequency and Risk Factors of Gout Flares in a Large Population-Based Cohort of Incident Gout | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 222 | Luise C, Trifiro` G, Mokhles M, Dieleman J, Soest E, Mazzaglia G, Herings R, Ross D & Sturkenboom M | Heterogeneity in Drug Utilization in Three European Countries: A Study of Dopamine Agonists | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 221 | Martin-Merino E, Fortuny J, Plana E, Rivero E & Garcia-Rodriguez L | Incidence of Diabetic Retinopathy and Diabetic Macular Edema in Newly Diagnosed Type 1 and Type 2 Diabetic Patients | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 220 | Gastanaga V, Wenstowrth C, Ganesh V, Hughes G, Xue F, Verma M, Dimery I & Zhao S | Incidence rates of edema in solid tumor cancer patients in the THIN Database | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |

| | | | | | |
|-----|---|--|--|----------------|------|
| 219 | Gastanaga V, Wentworth C, Hughes G, Xue F, Ganesh V, Verma M, Dimery I & Zhao S | Incidence Rates of Retinal Artery Thrombosis in Solid Tumor Cancer Patients in the THIN Database | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 218 | Zhou X, Murugesan S, Wentworth C, Bhullar H & Bate A | Lessons Learned from Mapping the THIN Database to the Observational Medical Outcome Partnership (OMOP) Common Data Model (CDM) | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 217 | Arana A, Wentworth CW, Rivero E, Plana E & Conde E | Lymphoma among patients with atopic dermatitis treated with topical corticosteroids (TCS) and/or topical calcineurin inhibitors (TCIs) | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 216 | Rhim A, Haynes K, Hardy J, Testani J & Yang Y | Maternal Use of Proton Pump Inhibitors (PPI) during Pregnancy Is Associated with an Increased Risk for Cardiac Birth Defects: Analysis of 208,951 Pregnancies from the THIN Database | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 215 | Marston L, Carpenter J, Walters K, Morris R, Nazareth I & Petersen I | Missing data in primary care databases; problems and a possible solution | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 214 | Petersen I, Gilbert R, Rudolfi A, Evans S & Nazareth I | Oral Antibiotic Prescribing in Pregnancy in United Kingdom | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 213 | Maguire A, Blak B & Haynes K | Parameterizing Risk-Benefit Models from Primary Care Data: Recurrence and Exacerbation | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 212 | Margolis DJ, Fanelli M, Hoffstad O & Lewis JD. | Potential Association Between the Oral Tetracycline Class of Antimicrobials Used to Treat Acne and Inflammatory Bowel Disease. | The American Journal of Gastroenterology | 105(12):2610-6 | 2010 |
| 211 | Blak B T, Hards M, Dattani H & Thompson M | Prescribing Behaviour of Smoking Cessation Products in the UK – A THIN Database Study | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 210 | Blak BT, Hards M, Thompson M & Dattani H | Prescribing behaviour of smoking cessation products in the UK – a THIN database study | 26th International Conference on Pharmacoepidemiology & Therapeutic Risk Management (ICPE) | Brighton, UK | 2010 |
| 209 | Raman S, Marshall S & Sturmer T | Prescription Patterns among Children with Attention Deficit Hyperactivity Disorder (ADHD) in the United Kingdom (UK) | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 208 | Schade R, Dieleman JP & Sturkenboom MC | Prescription trends for dopamine agonists in UK primary care - The influence of safety alerts concerning valvular heart disease | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 207 | Gonzalez-Perez A, Schlienger RG & Garcia Rodriguez A | Risk of acute pancreatitis in association with type 2 diabetes mellitus | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 206 | Trifiro' G, Mokhles M, Dieleman J, Soest E, Mazzaglia G, Herings R, Luise C, Ross D, Brusselle G, Colao A, Haverkamp W, Schade R, Camp G, Zanettini R & Sturkenboom M | Risk of Cardiac Valve Regurgitation with Dopamine Agonist Use in Parkinson's Disease and Hyperprolactinemia | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 205 | Trifiro G, Mokhles M, Dieleman J, Soest E, Mazzaglia G, Herings R, Luise C, Ross D, Brusselle G, Colao A, Haverkamp W, Schade R, Camp G, Zanettini R & Sturkenboom M | Risk of heart failure with dopamine agonist use in Parkinson's disease | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 204 | Kimberly Forde K, Kevin Haynes K, James Lewis J & LoRe V | Risk of Myocardial Infarction in UK Primary Care Patients with Hepatitis C Virus Infection | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 203 | Petersen I, Gilbert R, Evans S, Marston L & Nazareth I | SSRI and Risk of Congenital Cardiac Abnormalities | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |

| | | | | | |
|-----|---|--|---|-----------------------------------|------|
| 202 | Raman S, Marshall S & Sturmer T | Stimulant Prescription and Risk of Injury among Children with Attention Deficit Hyperactivity Disorder (ADHD) in the United Kingdom (UK): Preliminary Results | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 201 | Serumaga B, Avery A & Elliott R | The Impact of Prescribing Guidelines on Physician Selection of Drug Therapy for Newly Diagnosed Patients with Hypertension in the UK | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 200 | Rubino A, Lapi F, Nicotra F, Zambon A, Scotti L, Thompson M, Vannacci A, Corrao G & Mugelli A | Use of serotonergic medications and cardiac valvulopathy: A nested case-control study in the Health Improvement Network (THIN) Database. | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 199 | Meropol S & Metlay J | Validating Pneumonia Hospitalizations in the THIN Database | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 198 | Seminara N, K Abuabara K, Shin D, Kimme S, Margolis D, Atroxel A & Gelfand J | Validation of Psoriasis and Its Prevalence, Treatment, and Severity in the Health Improvement Network (THIN) | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 197 | Denburg M, Haynes K, Shults J, Lewis J & Leonard M | Validation of the Health Improvement Network (THIN) Database for Epidemiologic Studies of Chronic Kidney Disease | Pharmacoepidemiology and Drug Safety | (19): S1-S347 | 2010 |
| 196 | Currie CJ, Gale EA & Poole CD | Estimation of primary care treatment costs and treatment efficacy for people with Type 1 and Type 2 diabetes in the United Kingdom from 1997 to 2007 | Diabetic Medicine | 27(8):938-48. | 2010 |
| 195 | Munson JC, Kreider M, Chen Z, Christie JD & Kimmel SE | Factors associated with the use of corticosteroids in the initial management of idiopathic pulmonary fibrosis. | Pharmacoepidemiology and Drug Safety | 19(7):756-62 | 2010 |
| 194 | Gribbin J, Hubbard R, Gladman JR, Smith C & Lewis S | Risk of falls associated with antihypertensive medication: population-based case-control study | Age and Ageing | 39(5):592-7 | 2010 |
| 193 | Arana A, Wentworth CE, Ayuso-Mateos JL & Arellano FM | Suicide-related events in patients treated with antiepileptic drugs | The New England Journal of Medicine | 363: 542-551 | 2010 |
| 192 | Rait G, Walters K, Bottomley C, Petersen I, Iliffe S & Nazareth I | Survival of people with clinical diagnosis of dementia in primary care: cohort study | British Medical Journal (BMJ) | 341:c3584. doi: 10.1136/bmj.c3584 | 2010 |
| 191 | Langley TE, Szatkowski L, Gibson J, Huang Y, McNeill A, Coleman T & Lewis S | Validation of The Health Improvement Network (THIN) primary care database for monitoring prescriptions for smoking cessation medications. | Pharmacoepidemiology and Drug Safety | 19(6):586-90 | 2010 |
| 190 | Blak BT, Wilson K, Metcalfe M, Maguire A & Hards M | Evaluation of varenicline as an aid to smoking cessation in UK general practice – a THIN database study | UK National Smoking Cessation Conference (UKNSCC) | Glasgow, Scotland | 2010 |
| 189 | Hippisley-Cox J & Coupland C | Predicting the risk of Chronic Kidney Disease in Men and Women in England and Wales: prospective derivation and external validation of the QKidneyScores | BMC Family Practice | 21;11:49 | 2010 |
| 188 | Hardoon SL, Whincup PH, Petersen I, Capewell S & Morris RW | Trends in longer-term survival following an acute myocardial infarction and prescribing of evidenced-based medications in primary care in the UK from 1991: a longitudinal population-based study. | Journal of Epidemiology & Community Health | 65(9):770-4 | 2010 |
| 187 | Ruigomez A, Martin-Merino E & Rodriguez LAG | Validation of ischemic cerebrovascular diagnoses in the health improvement network (THIN) | Pharmacoepidemiology and Drug Safety | 19(6):579-585 | 2010 |
| 186 | Collins GS & Altman DG | An independent and external validation of QRISK2 cardiovascular disease risk score: a prospective open cohort study | British Medical Journal (BMJ) | 340:c2442. doi: 10.1136/bmj.c2442 | 2010 |

| | | | | | |
|-----|--|--|--|--|------|
| 185 | González-Pérez A, Aponte Z, Vidaurre CF & Rodríguez LA | Anaphylaxis epidemiology in patients with and patients without asthma: a United Kingdom database review. | Journal of Allergy and Clinical Immunology | 125(5):1098-1104 | 2010 |
| 184 | Martín-Merino E, Ruigómez A, García Rodríguez LA, Wallander MA & Johansson S | Depression and treatment with antidepressants are associated with the development of gastro-oesophageal reflux disease. | Alimentary Pharmacology & Therapeutics | 31(10):1132-40 | 2010 |
| 183 | Blak BT, Hards M, Thompson M & Dattani H | Impact of the UK public smoking ban on the prescribing of smoking cessation therapy in primary care – a THIN database study | 15th annual international meeting (ISPOR) | Atlanta, USA | 2010 |
| 182 | Blak BT, Wilson K, Metcalfe M, Maguire A & Hards M | Evaluation of varenicline as an aid to smoking cessation in UK general practice - a THIN database study | Current Medical Research & Opinion | 26(4):861–870 | 2010 |
| 181 | Swedo SE, Schrag A, Gilbert R, Giovannoni G, Robertson MM, Metcalfe C, Ben-Shlomo Y & Gilbert DL | Streptococcal infection, Tourette syndrome, and OCD: is there a connection? | Neurology | 74(17):1397-8 | 2010 |
| 180 | Cooper C, Steinbuch M, Stevenson R, Miday R & Watts NB | The epidemiology of osteonecrosis: findings from the GPRD and THIN databases in the UK | Osteoporosis International | (4):569-77 | 2010 |
| 179 | García Rodríguez LA & Ruigómez A | Case validation in research using large databases | British Journal of General Practice | 60(572):160-1 | 2010 |
| 178 | Sladkevicius E, Nagy E, Lack G & Guest JF | Resource implications and budget impact of managing cow milk allergy in the UK | Journal of Medical Economics | 13(1):119-28 | 2010 |
| 177 | Blak BT, Wilson K, Metcalfe M, Maguire A & Hards M | Evaluation of varenicline as an aid to smoking cessation in UK general practice – a THIN database study | Society for Research on Nicotine and Tobacco | Baltimore, USA | 2010 |
| 176 | Ban L, Gibson JE, West J & Tata LJ | Association between perinatal depression in mothers and the risk of childhood infections in offspring: a population-based cohort study | BMC Public Health | 31(10):799 | 2010 |
| 175 | Hall G, McMahon A, Dain M & Home PD | Duration of first insulin therapy in previously uncontrolled type 2 diabetes: comparison of insulins glargine, detemir and NPH and association with glycemic control | Value in Health | 13(7):A298 | 2010 |
| 174 | Ruigómez A, Wallander MA, Lundborg P, Johansson S & Rodriguez LA | Gastroesophageal reflux disease in children and adolescents in primary care | Scandinavian Journal of Gastroenterology | 45(2):139-46 | 2010 |
| 173 | Blak BT, Lee J, Hards M & Sedani T | Has the Quality and Outcomes Framework Influenced Primary Care Data Recording? | Value in Health | 13(7):A416 | 2010 |
| 172 | Blak BT, Lee J, Hards M & Sedani T | Has the Quality and Outcomes Framework influenced primary care data recording? | ISPOR | | 2010 |
| 171 | Blak BT, Hards M & Lee J | How do THIN death data compare to national figures for each UK country? | ISPOR | | 2010 |
| 170 | Blak BT, Hards & Lee J | How do THIN Death Data Compare to National Figures for each UK Country? | Value in Health | 13 (7):A331 | 2010 |
| 169 | Blak BT, Hards M, Thompson M & Dattani H | Impact of the UK public smoking ban on the prescribing of smoking cessation therapy in primary care – a THIN database study | Value in Health | 13(3):A203 | 2010 |
| 168 | Evans ML, Sharplin P, Owens DR, Chamberlain GH, Longman AJ & McEwan P | Insulin usage in type 2 diabetes mellitus patients in UK clinical practice: a retrospective cohort-based analysis using the THIN database | British Journal of Diabetes & Vascular Disease | 10: 178: DOI: 10.1177/1474651410370659 | 2010 |

| | | | | | |
|-----|---|---|--|-------------------|------|
| 167 | Taylor RR & Guest JF | The cost-effectiveness of macrogol 3350 compared to lactulose in the treatment of adults suffering from chronic constipation in the UK. | Alimentary Pharmacology & Therapeutics | 31(2):302-12 | 2010 |
| 166 | Donaldson GC, Hurst JR, Smith CJ, Hubbard RB & Wedzicha JA | Increased risk of Myocardial Infarction and Stroke following Exacerbation of Chronic Obstructive Pulmonary Disease | Chest | 137(5):1091-7 | 2009 |
| 165 | Ward K & Hubbard R | Is adult height related to the risk of having Chronic Obstructive Pulmonary Disease (COPD)? | Journal of Epidemiology & Community Health | 65(3):226-9 | 2009 |
| 164 | Sjöberg Bexelius T, García Rodríguez LA & Lindblad M | Use of angiotensin II receptor blockers and the risk of acute pancreatitis: a nested case-control study. | Pancreatology | 9(6):786-92 | 2009 |
| 163 | Rodríguez LA, Ruigómez A, Wallander MA & Johansson S | Acid-suppressive drugs and community-acquired pneumonia. | Epidemiology | 20(6):800-6 | 2009 |
| 162 | D Irwin, P Mernagh, S Thom pson, P Anderson & W Sawyer | Cost of Managing Neuropathic Pain with Pregabalin and Gabapentin: A UK Study | Poster presented at the RCGP Annual National Primary Care Conference | Glasgow, Scotland | 2009 |
| 161 | Ryan R P, McManus R J, Mant J, Macleod J A & Hobbs R | Statins in heart failure: Retrospective cohort study using routine primary care data | Annals of Medicine (informa healthcare) | (11):1-7 | 2009 |
| 160 | Cai S, García Rodríguez LA, Massó-González EL & Hernández-Díaz S | Uncomplicated peptic ulcer in the UK: trends from 1997 to 2005 | Alimentary Pharmacology & Therapeutics | 30(10):1039-48 | 2009 |
| 159 | Blak BT & Thompson M | How does The Health Improvement Network (THIN) data on prevalence of chronic diseases compare with national figures? | 12th Annual European Congress (ISPOR) | Paris, France | 2009 |
| 158 | Myles PR, Hubbard RB, Gibson JE, Pogson Z, Smith CJ & McKeever TM | Pneumonia mortality in a UK general practice population cohort | European Journal of Public Health | 19(5):521-6 | 2009 |
| 157 | Meropol SB, Chen Z & Metlay JP | Reduced antibiotic prescribing for acute respiratory infections in adults and children. | British Journal of General Practice | 59(567):e321-8 | 2009 |
| 156 | Schrag A, Gilbert R, Giovannoni G, Robertson M M, Metcalfe C & Ben-Shlomo Y | Streptococcal infection, Tourette syndrome, and OCD | Neurology | (73):1256-1263 | 2009 |
| 155 | Bhayat F, Das-Gupta E, Smith C & Hubbard R | NSAID use and risk of leukaemia: a population-based case-control study | Pharmacoepidemiology and Drug Safety | 18(9):833-6 | 2009 |
| 154 | Currie CJ, Poole CD & Gale EA | The influence of glucose-lowering therapies on cancer risk in type 2 diabetes | Diabetologia | 52(9):1766-77 | 2009 |
| 153 | LoRe V, Haynes K, Forde KA, Localio AR, Schinnar R & Lewis JD | Validity of The Health Improvement Network (THIN) for epidemiologic studies of hepatitis C virus infection | Pharmacoepidemiology and Drug Safety | 18(9):807-14 | 2009 |
| 152 | Langan SM, Hubbard R, Fleming K & West J | A population-based study of acute medical conditions associated with bullous pemphigoid. | British Journal of Dermatology | 161(5):1149-52 | 2009 |
| 151 | Marfatia A, Maneno MK, Li J & Jones JK | Longitudinal Assessment of Smoking in The Health Improvement Network (THIN) Database | 25th Intl Conference on Pharmacoepidemiology & Therapeutic Risk Management | Rhode Island, USA | 2009 |
| 150 | Collins GS & Altman DG | An independent external validation and evaluation of QRISK cardiovascular risk prediction: a prospective open cohort study | British Medical Journal (BMJ) | 339:b2584 | 2009 |
| 149 | Petersen I, Gilbert R, Ridolfi A, Evans S & Nazareth I | Antibiotic prescribing during pregnancy in primary care | The Society for Academic Primary Care (SAPC) | | 2009 |
| 148 | Szatkowski L, Coleman T, Lewis S & McNeil A | Can national smoking prevalence be monitored using data from primary care medical records? | The Society for Academic Primary Care (SAPC) | | 2009 |
| 147 | Gribbin J, Hubbard R, Smith C, Gladman J & Lewis S | Incidence and mortality of falls amongst older people in primary care in the United Kingdom | QJM: An International Journal of Medicine | 102(7):477-83 | 2009 |

| | | | | | |
|-----|---|---|--|-----------------|------|
| 146 | Schembri S, Morant S, Winter JH & MacDonald TM | Influenza but not pneumococcal vaccination protects against all-cause mortality in patients with COPD | Thorax BMJ | 64(7):567-72 | 2009 |
| 145 | Bhayat F, Das-Gupta E, Smith C, McKeever T & Hubbard R | The incidence of and mortality from leukaemias in the UK: a general population-based study | BMC Cancer | 9(1):252 | 2009 |
| 144 | Gribbin J, Hubbard R & Smith C | Role of diabetes mellitus and gastro-oesophageal reflux in the aetiology of idiopathic pulmonary fibrosis | Respiratory Medicine | 103(6):927-31 | 2009 |
| 143 | Blak BT, Hards M, Maguire A & Schwalm MS | Can the relationship between age and cholesterol levels indicate effectiveness of lipid lowering therapy in primary care? France, Germany, Italy and UK | ISPOR | Orlando, USA | 2009 |
| 142 | Haynes K, Forde KA, Schinnar R, Wong P, Strom BL & Lewis JD | Cancer incidence in The Health Improvement Network | Pharmacoepidemiology and Drug Safety | 18(8):730-736 | 2009 |
| 141 | Arellano FM, Arana A, Wentworth CE, Fernández-Vidaurre C, Schlienger RG & Conde E | Lymphoma among patients with atopic dermatitis and/or treated with topical immunosuppressants in the United Kingdom | Journal of Allergy and Clinical Immunology | 123(5):1111-6 | 2009 |
| 140 | Myles PR, Hubbard RB, Gibson JE, Pogson Z, Smith CJ & McKeever TM | The impact of statins, ace inhibitors and gastric acid suppressants on pneumonia mortality in a UK general practice population cohort | Pharmacoepidemiology and Drug Safety | 18(8):697-703 | 2009 |
| 139 | Myles PR, McKeever TM, Pogson Z, Smith CJ & Hubbard RB | The incidence of pneumonia using data from a computerized general practice database | Epidemiology and Infection | 137(5):709-16 | 2009 |
| 138 | Morant S, Bloomfield G, Vats V & Chapple C | Increased Sexual Dysfunction in Men with Storage and Voiding Lower Urinary Tract Symptoms | The Journal of Sexual Medicine | 6:1103-1110 | 2009 |
| 137 | Myles PR, Hubbard RB, McKeever TM, Pogson Z, Smith CJ & Gibson JE | Risk of community-acquired pneumonia and the use of statins, ace inhibitors and gastric acid suppressants: a population-based case-control study | Pharmacoepidemiology and Drug Safety | 18(4):269-75 | 2009 |
| 136 | Raine R, Wong W, Ambler G, Hardoon S, Petersen I, Morris R, Bartley M & Blane D | Sociodemographic variations in the contribution of secondary drug prevention to stroke survival at middle and older ages: cohort study | British Medical Journal (BMJ) | 338:b2095 | 2009 |
| 135 | Hamilton W, Lancashire R, Sharp D, Peters TJ, Cheng K & Marshall T | The risk of colorectal cancer with symptoms at different ages and between the sexes: a case-control study | BMC Medicine | 17(7):17 | 2009 |
| 134 | González EL, Johansson S, Wallander MA & Rodríguez LA. | Trends in the prevalence and incidence of diabetes in the UK: 1996-2005 | Journal of Epidemiology & Community Health | 63(4):332-6 | 2009 |
| 133 | Sharplin P, Gordon J, Peters JR, Tetlow AP, Longman AJ & McEwan P | Switching from premixed insulin to glargine-based insulin regimen improves glycaemic control in patients with type 1 or type 2 diabetes: a retrospective primary-care-based analysis | Cardiovascular Diabetology | 16(8):9 | 2009 |
| 132 | Hall G C | Validation of death and suicide recording on the THIN UK primary care database | Pharmacoepidemiology and Drug Safety | 18(2):120-31 | 2009 |
| 131 | Smith HT, Blak BT, Hards M & Arrelano J | "The "who, how and then what..." of insulin initiation in UK primary care" | Value in Health | 12(7):A412 | 2009 |
| 130 | Poole CD, Conway P & Currie CJ | An evaluation of the association between C-reactive protein, the change in C-reactive protein over one year, and all-cause mortality in chronic immune-mediated inflammatory disease managed in UK general practice | Rheumatology (Oxford) | 48(1):78-82 | 2009 |
| 129 | Blak BT, Maguire A, Schwalm MS, Hards M | Can the relationship between age and cholesterol levels indicate effectiveness of lipid lowering therapy in primary care? France, Germany, Italy and UK | Value in Health | 12(3):A142-A143 | 2009 |

| | | | | | |
|-----|---|--|--|-----------------|------|
| 128 | Smeeth L, Douglas I, Hall A J, Hubbard R & Evans S | Effect of statins on a wide range of health outcomes: a cohort study validated by comparison with randomized trials | British Journal of Clinical Pharmacology | 67(1):99-109 | 2009 |
| 127 | Blak BT & Thompson M | How does The Health Improvement Network (THIN) data on prevalences on chronic diseases compare with national figures? | Value in Health | 12(7):A253 | 2009 |
| 126 | Sharplin P, Gordon J, Peters J R, Tetlow A P, Longman A J & McEwan P | Improved glycaemic control by switching from insulin NPH to insulin glargine: a retrospective observational study | Cardiovascular Diabetology | 19(8)3 | 2009 |
| 125 | Blak BT, Mullins CD, Shaya FT, Simoni-Wastila L, Cooke CE & Weir MR | Prescribing Trends and Drug Budget Impact of the ARBs in the UK | Value in Health | (12):302-308 | 2009 |
| 124 | Rait G, Walters K, Griffin M, Buszewicz M, Petersen I & Nazareth I | Recent trends in the incidence of recorded depression in primary care | The British Journal of Psychiatry | (195):520–524 | 2009 |
| 123 | Maguire A, Douglas I & Blak BT | The application of discrete even simulation to quantitative risk benefit analysis | Value in Health | 12(7):A453 | 2009 |
| 122 | Bulugahapitiya U, Siyambalapitiya S, Sithole J, Fernando DJ & Idris I | The clinical impact of identifying metabolic syndrome in patients with diabetes: a cross-sectional study | Diabetes and Vascular Disease Research | 6(1):21-4 | 2009 |
| 121 | Maguire A, Blak B T & Thompson M | The importance of defining periods of complete mortality reporting for research using automated data from primary care | Pharmacoepidemiology and Drug Safety | 18:76–83 | 2009 |
| 120 | Gibson JE, Hubbard RB, Smith CJ, Tata LJ, Britton JR & Fogarty AW | The Use of Self-Controlled Analytical Techniques to Assess the Temporal Association Between use of Prescription Medications and the Risk of Motor Vehicle Crashes | American Journal of Epidemiology | 169:761-8 | 2009 |
| 119 | Gopal M, Haynes K, Bellamy SL, & Arya LA | Discontinuation Rates of Anticholinergic Medications Used for the Treatment of Lower Urinary Tract Symptoms | American College of Obstetricians and Gynecologists | 112(6) | 2008 |
| 118 | Keenan HT, Hall GC & Marshall SW | Early head injury and attention deficit hyperactivity disorder: retrospective cohort study | British Medical Journal (BMJ) | 337(a1984) | 2008 |
| 117 | Tata TJ, Lewis SA, McKeever TM, Smith CJ, Doyle P, Smeeth L, Gibson JE & Hubbard RB | Effect of maternal asthma, exacerbations and asthma medication use on congenital malformations in offspring: a UK population-based study | Thorax BMJ | 63(11):981-7 | 2008 |
| 116 | García Rodríguez LA, Tacconelli S & Patrignani P | Role of dose potency in the prediction of risk of myocardial infarction associated with nonsteroidal anti-inflammatory drugs in the general population | Journal of the American College of Cardiology | 52(20):1628-36 | 2008 |
| 115 | Poole CD, Conway P, Reynolds A & Currie CJ | The association between C-reactive protein and the likelihood of progression to joint replacement in people with rheumatoid arthritis: a retrospective observational study | BMC Musculoskelet Disorders | 4(9):146 | 2008 |
| 114 | Berger A, Sadosky A, Dukes E, Edelsberg J & Oster G | Use of pregabalin in patients with painful neuropathic disorders under the care of general practitioners in the U.K | Pain Practice | 9(1):18-34 | 2008 |
| 113 | Massó González EL & García Rodríguez LA | Proton pump inhibitors reduce the long-term risk of recurrent upper gastrointestinal bleeding: an observational study | Alimentary Pharmacology & Therapeutics | 28(5):629-37 | 2008 |
| 112 | Hubbard RB, Smith C, Le Jeune I, Gribbin J & Fogarty AW | The association between idiopathic pulmonary fibrosis and vascular disease: a population-based study | American Journal of Respiratory and Critical Care Medicine | 178(12):1257-61 | 2008 |
| 111 | Bulugahapitiya U, Siyambalapitiya S, Sithole J, Fernando DJ & Idris I | Age threshold for routine aspirin treatment to prevent cardiovascular disease in patients without diabetes | Heart | 94(11):1429-32 | 2008 |
| 110 | Morgan O, Griffiths C & Majeed A | Antidepressant prescribing and changes in antidepressant poisoning mortality and suicide in England, 1993–2004 | Journal of Public Health | (30):60 - 68 | 2008 |

| | | | | | |
|-----|---|---|--|---------------------|------|
| 109 | Blak BT & Maguire A | Influence of affluence: the impact of social deprivation on mortality rates in THIN, a UK primary care database | 24th International Conference on Pharmacoepidemiology & Therapeutic Risk Management (ISPE) | Copenhagen, Denmark | 2008 |
| 108 | Hippisley-Cox J, Coupland C, Vinogradova Y, Robson J & Brindle P | Performance of the QRISK cardiovascular risk prediction algorithm in an independent UK sample of patients from general practice: a validation study | Heart | (94):34-39 | 2008 |
| 107 | Smith CJP, Gribbin J, Challen KB & Hubbard RB | The impact of the 2004 NICE guideline and 2003 General Medical Services contract on COPD in primary care in the UK | QJM: An International Journal of Medicine | 101(2):145-153 | 2008 |
| 106 | Maguire A, Blak BT, Thompson M & Hall G | The importance of defining periods of complete mortality reporting in automated patient databases | 24th International Conference on Pharmacoepidemiology & Therapeutic Risk Management (ISPE) | Copenhagen, Denmark | 2008 |
| 105 | Petersen I, Man S L, Evans S, Gilbert R & Nazareth I | Antidepressant prescribing during pregnancy | The Society For Academic Primary Care, 37th Annual Scientific Meeting | Galway, Ireland | 2008 |
| 104 | Margolis D J, Hoffstad O & Strom BL | Association between serious ischemic cardiac outcomes and medications used to treat diabetes | Pharmacoepidemiology and Drug Safety | 17(8):753-759 | 2008 |
| 103 | Langan SM, Smeeth L, Hubbard R, Fleming KM, Smith C J & West J | Bullous pemphigoid and pemphigus vulgaris -incidence and mortality in the UK: population based cohort study | British Medical Journal (BMJ) | 337(a180) | 2008 |
| 102 | Haynes K, Heitjan D, Kanetsky P & Hennessy S | Declining public health burden of digoxin toxicity from 1991 to 2004 | Clinical Pharmacology & Therapeutics | 84(1):90-4 | 2008 |
| 101 | Thompson R, O'Regan C, Morant S, Phillips B & Ong S | Measurement of baseline total cholesterol: new data from The Health Improvement Network (THIN) database | Primary Care Cardiovascular Journal | 1(2):107-111 | 2008 |
| 100 | Rait G, Walters K, Bottomley C, Nazareth I, Petersen I & Liffie S | Survival following diagnosis of dementia in primary care | The Society For Academic Primary Care, 37th Annual Scientific Meeting | Galway, Ireland | 2008 |
| 99 | Hamilton W, Lancashire R, Sharp D, Peters TJ, Cheng K & Marshall T | Symptomatic colorectal cancer: risk differences across age and between the sexes | Annals of Oncology | | 2008 |
| 98 | Hamilton W, Lancashire R, Sharp D, Peters T J, Cheng K & Marshall T | The importance of anaemia in diagnosing colorectal cancer: a case-control study using electronic primary care records | British Journal of Cancer | 98(2):323 – 327 | 2008 |
| 97 | Dave S, Petersen I, Sherr L & Nazareth I | Using Primary care databases to assess incidence rates of parental depression | The Society For Academic Primary Care, 37th Annual Scientific Meeting | Galway, Ireland | 2008 |
| 96 | Hamilton WT, Lancashire R, Sharp D, Peters TJ, Cheng K & Marshall T | Validation of the CAPER score for diagnosis for colorectal cancer and comparison with NICE guidance | The Society For Academic Primary Care, 37th Annual Scientific Meeting | Galway, Ireland | 2008 |
| 95 | Zhang H, García Rodríguez LA & Hernández-Díaz S. | Antibiotic use and the risk of lung cancer | Cancer Epidemiology | 17(6):1308-15 | 2008 |
| 94 | Molokhia M, McKeigue P, Curcin V & Majeed A | Statin Induced Myopathy and Myalgia: Time Trend Analysis and Comparison of Risk Associated with Statin Class from 1991–2006 | Public Library of Science (PLoS one) | 3(6):e2522 | 2008 |
| 93 | Gao S, Juhaeri J & Dai WS | The incidence rate of seizures in relation to BMI in UK adults | Obesity | 16(9):2126-32 | 2008 |
| 92 | Guest J, Clegga J & Helter M | Cost-effectiveness of macrogol 4000 compared to lactulose in the treatment of chronic functional constipation in the UK | Current Medical Research & Opinion | 24(7):1841–1852 | 2008 |
| 91 | Morant SV, Reilly K, Bloomfield GA & Chapple C | Diagnosis and treatment of lower urinary tract symptoms suggestive of overactive bladder and bladder outlet obstruction among men in general practice in the UK | International Journal of Clinical Practice | 62(5):688–694 | 2008 |

| | | | | | |
|----|--|---|--|---------------|------|
| 90 | Morant et al | Master of nothing but knower of all | International Journal of Clinical Practice | 62(5):667-676 | 2008 |
| 89 | Hall GC, Carroll D & McQuay HJ. | Primary care incidence and treatment of four neuropathic pain conditions: a descriptive study, 2002-2005 | BMC Family Practice | 6(9):26 | 2008 |
| 88 | Vats V, Morant S, Chapple C & Kelleher C | Increased Sexual Dysfunction in Women With LUTS, Including OAB, in a UK General Practice Setting: Analyses From the THIN Database | 23rd Annual European Association of Urology Congress | Milan, Italy | 2008 |
| 87 | Harrison WN, Lancashire RJ & Marshall TP | Variation in recorded blood pressure terminal digit bias in general practice | Journal of Human Hypertension | 22(3):163-7 | 2008 |
| 86 | MacDonald TM & Morant SV | Prevalence and treatment of isolated and concurrent hypertension and hypercholesterolaemia in the United Kingdom | British Journal of Clinical Pharmacology | 65(5):775-786 | 2008 |
| 85 | Meropol S, Chen Z & Metlay J | Adverse events associated with pediatric antibiotic use | Pharmacoepidemiology and Drug Safety | 17(S273) | 2008 |
| 84 | Meropol S, Chen Z & Metlay J | Antibiotics for upper respiratory infections | Pharmacoepidemiology and Drug Safety | 17(S209-S210) | 2008 |
| 83 | Carrigan G, Khromava A & Holden W | Changes in the incidence of encephalitis in the UK | Pharmacoepidemiology and Drug Safety | 17(S253) | 2008 |
| 82 | Yuan Z, Weinstein R, Zhang J, Cheng M, Griffin G, Zolynas R, Lee M & Oppenheimer L | Clinical outcomes associated with anticoagulant therapies in patient with incident heart failure: A retrospective analysis of a population based database | Pharmacoepidemiology and Drug Safety | 17(S1-S294) | 2008 |
| 81 | Hall G | Death recording in a UK primary care database | Pharmacoepidemiology and Drug Safety | 17(S245-S246) | 2008 |
| 80 | Gulliford MC, Latinovic R & Charlton J | Diabetes Diagnosis, Resource Utilization, and Health Outcomes | American Journal of Managed Care | 14(1):32-8 | 2008 |
| 79 | Carnahan D, Haynes K, Sammel M, Holmes J & Turner B | Does an increase in serum creatinine after starting ACE inhibitors influence their discontinuation | Pharmacoepidemiology and Drug Safety | 17(S277) | 2008 |
| 78 | Blak B & Maguire A | Influence of affluence: The impact of social deprivation on mortality rates in THIN, a UK primary care database | Pharmacoepidemiology and Drug Safety | 17(S252) | 2008 |
| 77 | Schembri S, Morant S, Winter J & MacDonald T | Influenza and pneumococcal vaccination protect against all cause mortality in patients with chronic obstructive pulmonary diseases | Pharmacoepidemiology and Drug Safety | 17(S194) | 2008 |
| 76 | Arellano F, Arana A, Wentworth C, Conde E, Fernandez-Vidauure C & Schlienger R | Occurrence of Lymphoma among patients with atopic dermatitis and/or treated with topical immunosuppressants in the United Kingdom | Pharmacoepidemiology and Drug Safety | 17(S109) | 2008 |
| 75 | Maguire A & Schwalm MS | One question, several countries: understanding health systems and standardising recording habits in primary care databases in France, Germany, Italy and the UK | Pharmacoepidemiology and Drug Safety | 17(S279) | 2008 |
| 74 | Margulis A, Rodriguez L & Hernandez-Diaz S | Positive predictive value of computerised medical records for uncomplicated upper gastrointestinal ulcer | Pharmacoepidemiology and Drug Safety | 17(S126) | 2008 |
| 73 | Haynes K, Schinnar R, Wong P, Strom B & Lewis J | Recording of pancreatic, colorectal and lung cancer incidence in The Health Improvement Network (THIN) | Pharmacoepidemiology and Drug Safety | 17(S38) | 2008 |
| 72 | Tennis P, Roskell N, Andrews E & Rothman K | Risk factors for thiazolidinedione related peripheral edema | Pharmacoepidemiology and Drug Safety | 17(S118-S119) | 2008 |
| 71 | Laudati J, Motsko S & Jones J | Risk of chronic end stage renal disease after use of ACE inhibitors compared to other antihypertensives or no-antihypertensive use in a cohort of diabetic patients in the United Kingdom | Pharmacoepidemiology and Drug Safety | 17(S233) | 2008 |

| | | | | | |
|----|--|---|---|------------------|------|
| 70 | Maguire A, Blak B, Thompson M & Hall G | The importance of defining periods of complete mortality reporting in automated patient databases | Pharmacoepidemiology and Drug Safety | 17(S250) | 2008 |
| 69 | Arellano F, Conde E, Wentworth C, Schlienger R, Fernandez-Vidauure C & Arana A | Validation of cases of Lymphoma in THIN | Pharmacoepidemiology and Drug Safety | 17(S87-S88) | 2008 |
| 68 | Meal A, Leonardi-Bee J, Smith C, Hubbard R & Bath-Hextall F | Validation of THIN data for non-melanoma skin cancer | Quality in primary care journal | 16(1):49-52 | 2008 |
| 67 | Rubino A, McQuay L J, Gough SC, Kvasz M & Tennis P | Delayed initiation of subcutaneous insulin therapy after failure of oral glucose-lowering agents in patients with Type 2 diabetes: a population-based analysis in the UK | Diabetic Medicine | 24(12):1412-8 | 2007 |
| 66 | Le Jeune I, Gribbin J, West J, Smith C, Cullinan P & Hubbard R | The incidence of cancer in patients with idiopathic pulmonary fibrosis and sarcoidosis in the UK | Respiratory Medicine | 101(12):2534-40 | 2007 |
| 65 | Blak BT, Maguire A & Schwalm MS | Differences in cholesterol levels between statin and fibrate treated patients in French, Italian, and United Kingdom primary care | Value in Health | 10(6):A407 | 2007 |
| 64 | Bath-Hextall F, Leonardi-Bee J, Smith C, Meal A & Hubbard R | Trends in incidence of skin basal cell carcinom: Additional evidence from a UK primary care database study | International Journal of Cancer | 121(9):2105-8 | 2007 |
| 63 | Van Staa TP, Kanis JA, Geusens P, Boonen A, Leufkens HG & Cooper C | The cost-effectiveness of bisphosphonates in postmenopausal women based on individual long-term fracture risks | Value in Health | 10(5):348-57 | 2007 |
| 62 | Pujades-Rodriguez M, Smith CJ & Hubbard RB | Inhaled corticosteroids and the risk of fracture in chronic obstructive pulmonary disease | QJM: An International Journal of Medicine | 100(8):509-17 | 2007 |
| 61 | Gibson JE, Hubbard RB, Smith CJP, Tata LJ, Britton J & Forgarty AW | A general population based study of the impact of prescription-only medicines on the risk of road traffic accidents | Pharmacoepidemiology and Drug Safety | 16:S1-S274 (59) | 2007 |
| 60 | Douglas IJ, Smeeth & Evans SJ | A propensity score controlled cohort study to investigate the effects of Statins | Pharmacoepidemiology and Drug Safety | 16:S1-S274 (474) | 2007 |
| 59 | Jara M, Lanes SF, Wentworth C, May C & Kesten S | Comparative Safety of Long-Acting Inhaled Bronchodilators | Drug Safety | 30(12):1151-1160 | 2007 |
| 58 | Hall GC & McMahon AD | Comparative study of modified release alpha-blocker exposure in elderly patients with fractures | Pharmacoepidemiology and Drug Safety | 16(8):901-907 | 2007 |
| 57 | Tata LJ, Lewis SA, McKeever TM, Smith CJP, Doyle P, Smeeth L, Gibson JE & Hubbard RB | Congenital malformation in children born to women with asthma | Pharmacoepidemiology and Drug Safety | 16(S1-S274):271 | 2007 |
| 56 | Davies AR, Smeeth L & Grundy EM | Contribution of changes in incidence and mortality to trends in the prevalence of coronary heart disease in the UK: 1996 2005 | European Heart Journal | 28: 2142-2147 | 2007 |
| 55 | Hall G & McMahon A | The association of weight change with depression | Pharmacoepidemiology and Drug Safety | 16:S1-S220 (464) | 2007 |
| 54 | Currie CJ, Poole CD, Tetlow T, Holmes P & McEwan P | The outcome of care in people with type 1 and type 2 diabetes following switching to treatment with either insulin glargine or insulin detemir in routine general practice in the UK: a retrospective database analysis | Current Medical Research & Opinion | 23(1):33-9 | 2007 |
| 53 | Poole C, Tetlow D, McEwan T, Holmes P, Currie P & Craig J | The prescription cost of managing people with type 1 and type 2 diabetes following initiation of treatment with either insulin glargine or insulin detemir in routine general practice in the UK: a retrospective database analysis | Current Medical Research & Opinion | (8):41-48 | 2007 |
| 52 | Toh S, García Rodríguez LA & Hernández-Díaz S | Use of antidepressants and risk of lung cancer | Cancer Causes Control | 18(10):1055-64 | 2007 |

| | | | | | |
|----|---|--|---|----------------------|------|
| 51 | Tata L J, Lewis S A, McKeever T M, Smith C J, Doyle P, Smeeth L, West J & Hubbard RB | A comprehensive analysis of adverse obstetric and pediatric complications in women with asthma | American Journal of Respiratory and Critical Care Medicine | 175(10):991-7 | 2007 |
| 50 | Kell P D, Hvidsten K, Morant S V, Harnett J P & Bridge S | Factors that predict changing the type of phosphodiesterase type 5 inhibitor medication among men in the UK | British Journal of Urology | 99(4):860-3 | 2007 |
| 49 | Tata L J, Hubbard R B, McKeever T M, Smith C J P, Doyle P, Smeeth L, West J & Lewis S A | Fertility Rates in Women with Asthma, Eczema, and Hay Fever: A General Population-based Cohort Study | American Journal of Epidemiology | 165(9):1023-30 | 2007 |
| 48 | Coleman T, Lewis S, Hubbard R & Smith C | Impact of contractual financial incentives on the ascertainment and management of smoking in primary care | Addiction | 102(5):803-8 | 2007 |
| 47 | Ryan R, McManus RJ, Mant J, Macleod J & Hobbs R | The prescribing of statins and mortality following a diagnosis of heart failure | Annual Research Conference - West Midlands Deanery, Department of Primary Care & General Practice | Birmingham, UK | 2007 |
| 46 | Solaymani-Dodaran M, West J, Smith C & Hubbard R | Extrinsic Allergic Alveolitis: Incidence and mortality in the general population | QJM: An International Journal of Medicine | 100(4):233-7 | 2007 |
| 45 | Hernandez-Diaz S & Garcia Rodriguez LA | Non-steroidal anti-inflammatory drugs and risk of lung cancer | International Journal of Cancer | 120:1565–1572 | 2007 |
| 44 | Garcia Rodriguez LA & Barreales Tolosa L | Risk of upper gastrointestinal complications among users of traditional NSAIDs and COXIBs in the general population | Gastroenterology | 132(2): 498-506 | 2007 |
| 43 | Lewis JD, Schinnar R, Bilker WB, Wang X & Strom BL | Validation studies of The Health Improvement Network (THIN) database for pharmacoepidemiology research | Pharmacoepidemiology and Drug Safety | 16(4):393-401 | 2007 |
| 42 | Maguire A, Schwalm MS, Douglas I & Smeeth L | Classification of hyperlipidemia from automated primary care data in the UK and France | Pharmacoepidemiology and Drug Safety | 16:S1-S274 (424) | 2007 |
| 41 | Siyambalapitiya S, Bulugahapitiya U, Sithole J, Song S, Fernando D J S & Idris I | Combining Population Health and Baseline Risk Strategy by Determining an Age Cutoff for Initiating Statins in Patients With Diabetes | Diabetes Care | 30:2025-2029 | 2007 |
| 40 | Maguire A, Douglas, Smeeth L & Thompson M | Determinants of cholesterol and triglyceride recording in patients treated with lipid lowering therapy in UK primary care | Pharmacoepidemiology and Drug Safety | 16:S1-S274 (480) | 2007 |
| 39 | Blak BT, Maguire A & Schwalm MS | Differences in cholesterol levels between statin and fibrate treated patients in French, Italian and UK primary care | ISPOR | | 2007 |
| 38 | Maguire A, Douglas I & Smeeth L | Patterns of cholesterol levels in patients treated with lipid lowering drugs in UK primary care | Pharmacoepidemiology and Drug Safety | 16:S1-S274 (481) | 2007 |
| 37 | Cox D & Tennis P | Risk Factors for Peripheral Edema in Patients Treated With Rosiglitazone and Pioglitazone: a Retrospective Analysis Using the Health Improvement Network (THIN) Database | RTI Health Solutions - Report | | 2007 |
| 36 | Cooper SC, Graham P & Trudgill NJ | Risk factors for the development of oesophageal cancer in Barrett's oesophagus: a primary care cohort study | Gastroenterology | 132(4) Suppl 2: A420 | 2007 |
| 35 | Van Staa TP, Geusens P, Kanis JA, eufkens HGM, Gehlbach S & Cooper C | A simple clinical score for estimating the long-term risk of fracture in postmenopausal women | QJM: An International Journal of Medicine | 99(10):673-682 | 2006 |
| 34 | Blak BT, Weir MR, Simoni-Wastila L, Shaya FT, Cooke CE & Mullins CD | Antihypertensive drug budget impact of the Angiotensin II Receptor Blockers (ARB) in the UK | Value in Health | 9(6):A343 | 2006 |
| 33 | Blak BT, Weir MR, Simoni-Wastila L, Shaya FT, Cooke CE & Mullins CD | Antihypertensive Drug Budget Impact of the Angiotensin II Receptor Blockers (ARBs) in the UK | ISPOR | | 2006 |
| 32 | Haynes K, Thompson M, Dattani H, Bourke A & Hennessy S | Challenges in obtaining ethical approval for a pilot study to obtain biosamples in a UK medical records database | Pharmacoepidemiology and Drug Safety | 15:(S1):S137-138 | 2006 |

| | | | | | |
|----|---|--|---|----------------------|------|
| 31 | Jara MM, Lanes SF, Wentworth C, May C & Kesten S | Comparative safety of long-acting bronchodilators | Pharmacoepidemiology and Drug Safety | 15: (S1):S75 | 2006 |
| 30 | Price D, Morant SV, Horne R, Pettitt D & MacDonald T | Determinants of long acting bronchodilator adherence in UK COPD patients | 16th ERS Annual Congress | Munich, Germany | 2006 |
| 29 | Reilly K, Morant S, Bloomfield G & Chapple C | Diagnosis and Treatment of Lower Urinary Tract Symptoms, Overactive Bladder, and Benign Prostatic Hyperplasia in Men in UK General Practice | 36th Annual Meeting of the International Continence Society | | 2006 |
| 28 | Kell P, Hvidsten K, Morant S, Harnett J & Bridge S | Factors That Predict Switching Phosphodiesterase Type 5 Inhibitor Medication in the United Kingdom | The Journal of Sexual Medicine | 3(s3):199 | 2006 |
| 27 | Blak BT, Dattani H, Bourke A & Page M | Has the Quality and Outcomes Framework (QOF) Scheme Impacted upon General Practitioners' (GPs) Clinical Practice? | The National Institute of Clinical Excellence (NICE) conference | Birmingham, UK | 2006 |
| 26 | Gribbin J, Hubbard RB, Le Jeune I, Smith CJP, West J & Tata LJ | Incidence and mortality of idiopathic pulmonary fibrosis and sarcoidosis in the UK | Thorax BMJ | 61:980-985 | 2006 |
| 25 | Gulliford MC, Latinovic R, Charlton J, & Hughes RAC | Increased incidence of carpal tunnel syndrome up to 10 years before diagnosis of diabetes | Diabetes Care | 29(8): 1929-1930 | 2006 |
| 24 | Blak BT, Thompson M & Bourke A | National Representativeness and Data Quality of the Health Improvement Network (THIN) Database of Primary Care Information for Epidemiological Research | 9th Annual Conference of the UK Federation of Primary Care Research Organisations | Liverpool, UK | 2006 |
| 23 | Mortimer K, Tata LJ, Smith CJP, West J, Harrison TW, Tattersfield A & Hubbard R | Oral and inhaled corticosteroids and adrenal insufficiency: a case-control study | Thorax BMJ | 61(5): 405-8 | 2006 |
| 22 | Rubino A, McQuay L, Kvasz M & Tennis P | Population-Based Study of Time to Insulin After Failure of Oral Antidiabetic (OA) Therapy in Type 2 Diabetes (T2DM) in the UK | Annual meeting of the American Diabetes Association | Washington, USA | 2006 |
| 21 | Smeeth L, Cook C, Thomas S, Hall AJ, Hubbard R & Vallance P | Risk of deep vein thrombosis and pulmonary embolism after acute infection in a community setting | Lancet | 367: 1075-79 | 2006 |
| 20 | Gulliford MC, Charlton J & Latinovic R | Risk of Diabetes Associated With Prescribed Glucocorticoids in a Large Population | Diabetes Care | 29 (12): 2728-29 | 2006 |
| 19 | Rubino A, McQuay L, Kvasz M & Tennis P | Time to insulin after failure of oral antidiabetic therapy in type 2 diabetes: A UK population-based analysis | Pharmacoepidemiology and Drug Safety | 15: (S1):S51 | 2006 |
| 18 | Blak B T, Mullins CD, Simoni-Wastila L, Shaya FT, Cooke CE & Weir MR | Trends in Angiotensin II Receptor Blocker (ARB) prescribing among general practitioners in the UK | Value in Health | 9(6):A204 | 2006 |
| 17 | Arellano FM, Yood MU, Wentworth CE, Oliveria SA, Rivero E, Verma A & Rothman KJ | Use of cyclo-oxygenase 2 inhibitors (COX-2) and prescription non-steroidal anti-inflammatory drugs (NSAIDs) in UK and USA populations Implications for COX-2 cardiovascular profile | Pharmacoepidemiology and Drug Safety | 15:861-872 | 2006 |
| 16 | Lewis JD, Schinnar R, Bilker WB, Wang X & Strom BL | Validation studies of The Health Improvement Network (THIN) Database for pharmacoepidemiology research | Pharmacoepidemiology and Drug Safety | 15: (S1):S233-234 | 2006 |
| 15 | Alder J, Mayhew L, Moody S, Morris R & Shah R | The chronic disease burden - An analysis of health risks and health care usage | Cass Business School (City University) | ISBN 1 - 901615-87-1 | 2005 |
| 14 | Van Staa TP, Geusens P, Leufkens HGM & Cooper C | Persistence to bisphosphonate treatment in actual clinical practice | Bone | 36 (S2):S412 | 2005 |
| 13 | Arellano FM, Wentworth CE, May C, Verma A & Cid J | Patterns of discontinuation, Dose modification and switching among users of prescription Non-Steroidal Anti-Inflammatory Drugs (NSAID) including Cyclo-Oxygenase 2 Inhibitors (COX-2) in the UK population | Pharmacoepidemiology and Drug Safety | 14: S1: (003) | 2005 |

| | | | | | |
|----|---|---|--|---------------------|------|
| 12 | Kell P, Hvidsten K, Morant S, Harnett J & Bridge S | Factors that predict switching Phosphodiesterase type 5 Inhibitor medication in the United Kingdom | 8th Congress of the European Society for Sexual Medicine | Copenhagen, Denmark | 2005 |
| 11 | Tata LJ, Fortune PJ, Hubbard RB, Smeeth L, Hawkey CJ, Smith CJP, Whitaker HJ, Farrington CP, Card TR & West J | Does concurrent prescription of selective serotonin reuptake inhibitors and non-steroidal anti-inflammatory drugs substantially increase the risk of upper gastrointestinal bleeding? | Alimentary Pharmacology & Therapeutics | 22: 175-181 | 2005 |
| 10 | Hubbard R, Lewis SA, West J, Smith C, Godfrey C, Smeeth L, Farrington P & Britton J | Bupropion and the risk of sudden death: a self-controlled case-series analysis using The Health Improvement Network | Thorax BMJ | 60(10): 848-50 | 2005 |
| 9 | Van Staa TP, Geusens P, Boonen A, Leufkens HGM & Cooper C | An individualised model of cost-effectiveness of bisphosphonates in elderly women | Bone | 36 (S2):S138 | 2005 |
| 8 | Van staa TP, Geusens P, Pols HAP, De Laet C & Leufkens HGM | A simple score for estimating the long-term risk of fracture in patients using oral glucocorticoids | QJM: An International Journal of Medicine | 98:191-198 | 2005 |
| 7 | Hubbard R, Lewis S, Godfrey C, Smeeth L & Farrington P | Use of nicotine replacement therapy and the risk of acute myocardial infarction, stroke and death | Tobacco Control | 14(6): 416-421 | 2005 |
| 6 | Arellano FM, Wentworth CE, May C, Verma A, Rivero E & Rothman KJ | Use of prescription Cyclo-Oxygenase 2 inhibitors (COX-2) and Non COX-2 Non-Steroidal Anti-Inflammatory Drugs (NSAID) in a UK population | Pharmacoepidemiology and Drug Safety | 14: S141: (283) | 2005 |
| 5 | Arellano FM, Yood MU, Wentworth CE, Oliveria SA & Rothman KJ | Use of prescription Non-Steroidal Anti-Inflammatory Drugs (NSAID) Including Cyclo-Oxygenase 2 Inhibitors (COX-2) in a UK Population and a US Population | Pharmacoepidemiology and Drug Safety | 14: S190: (382) | 2005 |
| 4 | Hubbard R, Smith C, Smeeth L, Farrington P, West J, Lewis S, Godfrey C & Britton J | The Risk of Fits, Diagnosis of Epilepsy and Death after First Prescription for Bupropion: An Analysis of The Health Improvement Network (THIN) Database | Pharmacoepidemiology and Drug Safety | (13): S22 (S1-S334) | 2004 |
| 3 | Van Staa T P, Geusens P, Leufkens H, Gehlbach S & Cooper C | The estimation of the 5-year probability of fracture in a general population of elderly women | Journal of Bone and Mineral Research | 19 (S1);S292 | 2004 |
| 2 | Page M & Dattani H | Feasibility Study and Methodology to Establish a System to Improve Recording of Quality Indicators in Primary Care Data | Annual Scientific Meeting, "Moving from policy to reality-How was it for you?" | | 2004 |
| 1 | Bourke A, Dattani H & Robinson M | Feasibility Study and Methodology to create a Quality Evaluated Database of Primary Care Data | Informatics in Primary Care | 12(3):171-7 | 2004 |

